



Assessment of the Opinion of Women Older than 30 about the Risk of Overweight and Obesity

Naimov Dilshod Qayim o'g'li

Bukhara State Medical Institute, Uzbekistan, Bukhara

Summary: The article presents the results of assessing the state of women's health in relation to abdominal obesity and assessing the degree of cardiovascular risk with abdominal obesity, overweight and obesity, identified by the Quetelet index.

Keywords: metabolic syndrome, abdominal obesity, overweight, obesity.

Relevance

The problem of obesity in combination with various metabolic disorders and/or diseases is the focus of modern medical science and public health. The prevalence of obesity in the world is so great that it has become a global epidemic. With the spread of obesity across the planet, the associated severe somatic diseases multiply and worsen: type 2 diabetes mellitus (DM2), arterial hypertension (AH), coronary heart disease, oncological diseases, etc., which reduce the quality of life and increase the mortality rate among the working population (1,2).

Obesity leads to the formation of a number of diseases, disability, premature mortality (5.6). Currently, more than 320 thousand deaths from diseases associated with obesity are recorded in Europe. Obesity, primarily abdominal obesity, is a leading risk factor for CVD and non-insulin-dependent diabetes mellitus). There is evidence (7) that the waist-to-hip ratio (W/H) is an independent risk factor for CVD and mortality. At the same time, the WC/OB ratio is more related to CVD than body mass index (BMI). The first reports on the relationship between the nature of the deposition of fat mass and the risk of CVD and DM were made more than half a century ago (4,7).

Abdominal obesity is a more significant predictor of morbidity and mortality than BMI. This fact is indicated by studies by AH Kissebah and P. Bjorntorp, who showed that with the same BMI, abdominal obesity is accompanied by a higher risk of comorbidities than peripheral obesity (3). This, to a certain extent, can explain the greater severity of complications of obesity in men than in women (1).

Purpose of the study. To study the opinion of women on the degree of risk of overweight and obesity as a risk factor for cardiovascular diseases at the level of primary health care

Materials and methods. Representative sample of the unorganized population aged 20 to 69 years in the amount of 797 people (242 men and 555 women). Metabolic syndrome was diagnosed in the presence of abdominal obesity (waist circumference > 94 cm for men and > 80 cm for women). Overweight was calculated using the formula: weight (kg)/height (m)², ≥ 25, and IC levels ≥ 30 were taken as obese.

Results and discussions: When studying the degree of cardiovascular risk in women with abdominal obesity, BMI and obesity, among all examined women (table 1), less than half (43.84%) consider abdominal obesity to be a serious risk factor for cardiovascular disease (CVD) .

Table 1. Assessment of the predictive degree of risk of abdominal obesity among the female population

Number of observations			
Women's opinion	AO no	AO is	Total
Much	112	130	242
Not really	72	142	214
Does not increase	16	12	28
Other reasons	35	33	68
Total	235	317	552
In percents			
Women's opinion	AO no	AO is	Total
Much	47.66	41.01	43.84
Not really	30.64	44.79	38.77
Does not increase	6.81	3.79	5.07
Other reasons	14.89	10.41	14.89
Total	100.00	100.00	100.00

At the same time, every fifth woman (19.96%) does not consider abdominal obesity a risk factor for CVD. It should be noted that even among women suffering from AO, only 2/5 recognize obesity as a serious risk factor for CVD (41.01%), and 44.79% of the surveyed answered that obesity is not a very important risk factor. Every seventh woman (14.2%) with AO does not consider obesity a risk factor for CVD.

It is especially important that among women without AO there is a more critical attitude towards increased weight than among women with AO. If 41.01% of women with AO consider increased weight to be an important risk factor for CVD, then 47.66% of women with normal body weight adhere to this opinion. Moreover, the revealed differences are statistically significant ($p < 0.05$).

Next, we analyzed the risk assessment among women with BMI and obesity, identified on the basis of elevated Quetelet index (Table 2). It was found that only about half of women with obesity (52.94%) and 2/5 of women with BMI (%) consider obesity to be a serious risk factor for CVD.

Table 2. Estimation of the prognostic degree of risk among women with BMI and obesity, identified on the basis of the Ket indexle

Number of observations				
Women's opinion	Norm	BMI	Obesity	Total
Much	140	48	54	242
Not really	142	41	31	214
Does not increase	16	eight	four	28
Other reasons	38	17	13	68
Total	336	114	102	552
In percents				
Women's opinion	Norm	BMI	Obesity	Total
Much	41.67	42.11	52.94	43.84
Not really	42.26	35.96	30.39	38.77
Does not increase	4.76	7.02	3.92	5.07
Other reasons	11.31	14.91	12.75	12.32
Total	100.00	100.00	100.00	100.00

Almost every third woman with BMI (35.96%) and obesity (30.39%) does not believe that increased weight significantly increases cardiovascular risk. 21.93% of women with BMI and 16.67% of obese women do not consider their increased body weight to be a risk factor for CVD at all.

Conclusion. Thus, an unfavorable situation has developed among the female population regarding the assessment of their body weight and the significance of increased weight in the formation of cardiovascular risk.

Literature.

1. A.S. Ametov, T.Yu. Demidova, M.V. Stelmakh, T.I. Depuis. Relationship between obesity and type 2 diabetes mellitus in the light of evidence-based medicine // Journal for the continuing medical education of physicians. Endocrinology: news, opinions, training №2 2016— page 38
2. Bokarev I.N. metabolic syndrome. Clinical Medicine, No. 8, 2014.
3. Zhuraeva H.I. Kayumov L.Kh., Ubaidova D.S., Dzhabborov Zh.Zh. Relationship between myocardial infarction and metabolic syndrome// Electronic scientific journal "Biology and Integrative Medicine". - 2019. - No. 8. - S. 66 - 77.
4. Aguilar M, Bhuket T, Torres S, Liu B, Wong RJ. Prevalence of the metabolic syndrome in the United States, 2003-2012. JAMA. 2015 May 19;313(19):1973-4.
5. International Diabetes Federation: The IDF consensus worldwide definition of the metabolic syndrome, <http://www.idf.org/metabolic-syndrome>
6. Jaspinder K. A Comprehensive Review on Metabolic Syndrome. Cardiology Research and Practice Volume 2014, Article ID 943162, 21 pages
7. Jo'rayeva H. I, Soliyev A. U, Djumayev K. Sh, Khusenov O. N. Analysis Of The Assessment Of The Attitude To The State Of Its Health Among The Inorgonized Population//European Journal of Molecular & Clinical Medicine 2020.12. 5.3346-3352pages.
8. DK Naimov LEFT VENTRICULAR HYPERTROPHY: DIAGNOSIS IN 40-60-YEAR-OLD WOMEN WITH HYPERTENSION.
9. Naimov Dilshod. ASSESSMENT OF WOMEN'S OPINION ON THE RISK OF OVERWEIGHT AND OBESITY Vol. 1 no. 6 (2021): SYNERGY: JOURNAL OF ETHICS AND GOVERNANCE
10. Lawler PR, Filion KB, Eisenberg MJ Efficacy of exercise-based cardiac rehabilitation post-myocardial infarction: a systematic review and meta-analysis of randomized controlled trials. Am Heart J, 2011, vol. 162, pp. 571–584.
11. DK Naimov. (2022). MYOCARDITIS AGAINST THE BACKGROUND OF COVID-19: CLINICAL FEATURES AND DRUG TREATMENT. Journal of Integrated Education and Research, 1(1), 497–512.
5. DK Naimov MEDICAL THERAPY IN PATIENTS WITH ACUTE HEART FAILURE
12. Kalenchyts T.I., Rysevets A.V., Antanovich Zh.V. Perenosimost' fizicheskikh training u patsientov s infarktom miokarda na etape rannei statsionarnoi reabilitatsii [Tolerability of physical training in patients with myocardial infarction at the early inpatient rehabilitation stage]. Cardiology in Belarus, 2017, vol. 9, no. 3, pp. 466-470 (in Russian).
13. Naimov DK (2022). LEFT VENTRICULAR HYPERTROPHY: DIAGNOSIS IN 40-60-YEAR-OLD WOMEN WITH HYPERTENSION. European Journal of Interdisciplinary Research and Development, 4, 186–188. Retrieved from <http://www.ejird.journalspark.org/index.php/ejird/article/view/86>
14. Antanovich Zh.V., Kalenchyts TI, Rysevets AV Otsenka effektivnosti fizicheskikh treniro - vok u patsientov s infarktom miokarda v usloviyah statsionarnogo otdeleniya me- ditsinskoi reabilitatsii [Evaluation of the physical training effectiveness in patients with myocardial infarction at

inpatient department of medical rehabilitation]. *Cardiology in Belarus*, 2017, vol. 9, no. 3, pp. 392–395 (in Russian).

15. Zabolotnih II, Kantemirova RK *Kliniko-ekspertnaya diagnostika patologii vnutrennih organov: rukovodstvo dlya vrachei* [Clinical and expert diagnostics of the internal organs pathology: a manual for doctors]. St. Petersburg: SpetsLit, 2007, 190 p. (in Russian).
16. Smichek VB *Reabilitatsiya bol'nih i invalidov* [Rehabilitation of patients and invalids]. Moscow: MedLit, 2010, 560 p. (in Russian).
17. Naimov DK (2022). LEFT VENTRICULAR HYPERTROPHY: DIAGNOSIS IN 40-60-YEAR-OLD WOMEN WITH HYPERTENSION. *European Journal of Interdisciplinary Research and Development*, 4, 186–188.