



Assessment of the Quantitative Composition of Phytoplankton in Bukhara and Navoy Cities. Buxoro Va

Jumaeva Sh. B. ¹

¹ Bukhara State Medical Institute, Uzbekistan

Annotation: In conditions of low water under the influence of intensive anthropogenic impact in Uzbekistan, the rational use of water resources in the national economy is important. In this regard, continuous monitoring, assessment of the variability of the microbial, chemical composition, hydro biological parameters of water in reservoirs is of great importance.

Keywords: mineral, Bukhara region, Bacillariophyta, Chlorophyta, Cyanophyta, Dinophyta, Euglenophyta.

Changes in the chemical and mineral composition of water affect the microbial composition of water in surface (open) water bodies. Pathogenic microorganisms transmitted by water, adapting to these conditions, change their biological properties [7].

Due to the vital activity of aquatic organisms, the chemical composition of water is formed, thereby determining its quality. The permanent microflora of water bodies performs the function of a primary oxidant or reductant of pollutants entering the water body [4].

An important part of aquatic ecosystems is aquatic biota, represented by a mosaic of interconnected biocenoses that occupy all possible biotopes in streams and reservoirs: they inhabit the water column (plankton), the thickness and surface of the soil (benthos), overgrow the surface of solid rocky substrates washed by water or settling on the surface of macrophytes and in the turf of water moss (periphyton). Their species composition and structure are entirely determined by climatic and landscape conditions prevailing in drainage basins or directly in river corridors and coastal zones of lakes [1, 2, 3].

Consequently, biocenoses can be considered as information systems that characterize the condition of not only specific water bodies, but also the surrounding natural complexes on the territory of which they are located [8].

Phytoplankton are microscopic plant organisms that freely hover in the water column and carry out photosynthesis, and are one of the important elements of aquatic ecosystems involved in the formation of water quality and the productivity of a reservoir [9, 10, 11].

As is known, the study of phytoplankton occupies an important place in biomonitoring, since algae assimilate solar energy, accumulating it in the form of organic compounds in the process of photosynthesis, while releasing oxygen necessary for the respiration of algae themselves and other inhabitants of the reservoir [5]. The organic matter synthesized by them serves as a source of energy for heterotrophic organisms - bacteria, animals. Therefore, the properties of the phytoplankton link of an ecosystem determine its condition.

The abundance, biomass, taxonomic composition, and physiological activity of phytoplankton make it possible to draw conclusions about the well-being of the water body or its critical condition [8].

The purpose of this research work was to study and evaluate the qualitative and quantitative composition of phytoplankton in the studied areas of surface reservoirs of the Bukhara region.

Materials and research methods The composition, abundance, and distribution of phytoplankton over surface (open) water bodies of the region studied by us (Bukhara region of the Republic of Uzbekistan) are due to different hydrological, hydrochemical conditions and anthropogenic impact.

We have investigated the surface water bodies of the Bukhara region - Lake Tudakul and the Kuyumazar reservoir, which are used for household, drinking, cultural and irrigation purposes. The researches were carried out in the spring and summer of 2020. There were examined 16 samples taken from different places of these reservoirs.

In biological research, the most reliable and acceptable method for sampling phytoplankton from surface water is the bathometric method. Samples taken with a bathometer were used both for quantitative accounting and for the qualitative characterization of the sample taken.

In the researches carried out, phytoplankton samples were taken with a one-liter Ruttner bathometer: samples were poured 250 ml into 500 ml dishes, mixed (i.e., integral samples were taken).

For high-quality collection of phytoplankton, we used silk gas plankton net No. 76. For "soft" fixation of phytoplankton samples, Lugol's solution (until slightly yellow) was used, followed by the addition of 40% formalin (10 ml of 40% formalin is sufficient for 0.5 l of the sample).

It should be taken into account that high concentrations of the specified fixative cause deformation of algae and a change in the color of their pigment [8].

Phytoplankton samples were collected according to generally accepted algological methods [5, 8]; identification guides were used to identify the species composition of microalgae [10, 11].

The sample taken in a polyethylene bottle was fixed with 40% formalin and Lugol's solution, supplied with a label (sample number, date, water body, by whom it was taken, in the presence of whom the sample was taken). Standard horizons for water sampling were: 0 (surface); 0.5; 1.0; 2.5; 5 m.

The sediment method was used for phytoplankton samples under conditions. Then, according to the methods, the precipitate was brought to 100 ml.

In office conditions, the sedimentation method (sedimentation) was used to concentrate samples, because Planktonic cells settle at a speed of 1 cm in 3 hours [9], then the samples were settled in a dark place for 5-10 days, and then the filtrate was slowly sucked off with a siphon through a double layer of plankton net from silk gas No. 76 (this helps to preserve the fine structures of algae).

The compaction of the taken sample was carried out in 2 stages: from 0.5 L (500 ml) to 0.1 L (100 ml). Then, after secondary settling (no more than 5 days), the solution was sucked off again. Poor samples (winter-spring) were brought up to a volume of 10 ml (usually up to 20 ml), very rich (summer ones during the "blooming" blue-green period) - up to 50 ml, sometimes even up to 100 ml, in this case the secondary settling of the sample is not produced).

The quantitative processing of the material was carried out according to the generally accepted algological method in a Goryaev chamber with a volume of (0.001 mm³) or in a Fuchs-Rosenthal chamber with a volume of (3.2 mm³). Further research was carried out in the laboratory using a MEIJI light microscope.

The data obtained by counting phytoplankton were used to calculate their abundance, for which the conversion formula was used:

$N = n \times v1 / v2 \times W$, where,

N is the number of cells in 1 cm³ of water;

N is the number of cells in a 1 mm³ chamber;

v1 - Volume of sample concentrate (cm³);

v2 - Chamber volume (cm³);

W - The volume of the taken sample (cm³).

The biomass of the studied phytoplankton was determined by a generally accepted calculation method [3, 8].

In the course of research, the following qualitative and quantitative characteristics of phytoplankton were determined: species composition of phytoplankton; the number of species in each main group; taxonomic composition of phytoplankton; number and biomass of species and main groups of phytoplankton - number of cells $\times 10^3$.

Research results and discussion During the reconnaissance trip, samples of phytoplankton were taken, in which 75 species, varieties and forms of algae were found: diatoms (*Bacillariophyta*) - 34 species; green (*Chlorophyta*) - 20 species; blue-green (*Cyanophyta*) - 15 species; dinophyta(*Dinophyta*) - 5 species; Euglenophyta(*Euglenophyta*) - 1 species.

The taxonomic structure of phytoplankton in water bodies of the Bukhara region is presented in table. 1.

Table1 Taxonomic structure of phytoplankton in the studied areas of water bodies in Bukhara region

Таксон водорослей	LakeTudakul	Kuyumazarreservoir
<i>Bacillariophyta</i> (diatoms)	13	21
<i>Chlorophyta</i> (green)	14	6
<i>Cyanophyta</i> (blue-green)	9	6
<i>Dinophyta</i> (dinophytic)	5	-
<i>Euglenophyta</i> (euglena)	1	-
Total number of species	42	33

The dominant complex of phytoplankton communities in the studied areas of Lake Tudakul and the Kuyumazar reservoir was mainly represented by producers. Among them, diatoms, green and blue-green algae reach the greatest development and diversity, as well as dinophytic and euglenic algae with a low abundance (1-5 species).

Diatoms (*Bacillariophyta*) and green (*Chlorophyta*) algae in the samples from Lake Tudakul were detected more often than others - 13 (31.0%) and 14 (33.3%) species, respectively.

Blue-green algae (*Cyanophyta*) in phytoplankton samples of the studied areas of this reservoir are poorly represented, only 9 species, which amounted to 21.43% of the total number of species. The widespread planktonic colonial and filamentous forms of algae of the genera *Merismopedia*, *Microcystis*, *Gloeocapsa*, *Gomposphaeria* and species of the *Oscillatoriaceae* family prevailed.

The most indicative quantitative development of blue-green algae (*Cyanophyta*) was noted in a sample from Lake Tudakul. The number of blue-greens here in the sample was 6500.00×10^3 cells / l, and the biomass - 58.694 mg / l. The least development of blue-green algae was noted in a sample from the Kuyumazar reservoir: the number was 706.250×10^3 cells / l with a biomass of 11.150 mg / l (Table 2).

Table2 Quantitative development of phytoplankton in the studied areas of water bodies in Bukhara region (number of cells $\times 10^3$ / biomass mg / l)

Taxon	LakeTudakul	Kuyumazarreservoir
<i>Cyanophyta</i> (blue-green)	6500,00 / 58,694	706,250 / 11,150
<i>Bacillariophyta</i> (diatoms)	162,500 / 61,344	193,750 / 187,800
<i>Euglenophyta</i> (euglena)	12,500 / 11,150	0
<i>Dinophyta</i> (dinophytic)	150,625 / 211,125	0
<i>Chlorophyta</i> (green)	756,250 / 188,400	87,500 / 31,500
Total number of cells $\times 10^3$ cells / l / biomass, mg / l	17621,875 / 530,713	987,500 / 230,450

Note: in the numerator is the total number of phytoplankton (cells / l), in the denominator is the biomass of phytoplankton (mg / l).

In terms of taxonomic diversity, diatoms (*Bacillariophyta*) occupied a dominant position in phytoplankton samples (34 species, 45.33%) of the studied open water bodies.

In June phytoplankton samples from Lake Tudakul, planktonic freshwater-brackish-water forms of b-mesosaprobic species from the genera *Melosira*, *Cyclotella*, *Fragilaria*, *Synedra* prevailed. In the phytoplankton sample of the Kuyumazar reservoir, brackish-water (mesohalobes) b- and b- α -mesosaprobic forms of algae from the genera *Cocconeis*, *Achnanthes*, *Gyrosigma*, *Amphiprora*, *Navicula*, *Bacillaria*, *Nitzschia* prevailed.

With a single occurrence, diatoms (*Bacillariophyta*), due to the large cell size, made up a large biomass. The number of diatoms in the samples was $162,500 \times 10^3$ cells / l and $193,750 \times 10^3$ cells / l, and the biomass, respectively, was 61.344 mg / l and 187,800 mg / l.

Green algae (*Chlorophyta*) in the samples of the investigated water bodies are moderately represented - 20 species or 26.67%, forms and varieties, which are mainly represented by widespread b-mesosaprobic species from the genera *Ankistrodesmus*, *Oocystis*, *Chlorella*, *Chlamidomonas*, *Scenedesmus*, *Cosmarium*. The number of green algae in phytoplankton samples was 756.250×10^3 cells / l and 87.500×10^3 cells / l, and the biomass was 188.400 mg / l and 31.500 mg / l, respectively.

In phytoplankton samples from Lake Tudakul, there was a good development of dinophytic algae - *Dinophyta* (5 species), which are mainly represented by the genera *Glenodinium*, *Peridinium*, and eugleicalgae - *Thalassomonas*, with a single occurrence. The number of dinophytic algae was 150.625×10^3 cells / l, and the biomass was 211.125 mg / l. At the same time, dinophytic and eugleic algae were not found in water samples from the Kuyumazar reservoir.

It should also be noted that representatives of rotifers (genera *Rotaria*, *Colurella*, *Lecane*, *Chilodonella*) and fungi (*Mucota spp.*) were found in phytoplankton samples from water bodies.

The obtained hydrobiological information of the studied water bodies of the Bukhara region indicates that anthropogenic factors, in particular the pollution of open water bodies, cause changes in the composition, structure and ecological condition of aquatic biocenoses of different depths, expressed in the change of dominant complexes of organisms, simplification of the ecological structure, the appearance in the composition dominants of highly advanced species.

Analysis of the ecological characteristics found in the phytoplankton of algae indicates that under the conditions of modern water salinity in the studied water bodies, the species composition of summer phytoplankton was mainly freshwater-brackish water b-, b- and α -mesosaprobic, brackish-water forms of algae. The increased abundance of phytoplankton was created mainly by representatives of colonial and filamentous blue-green algae of the *Oscillatoriaceae* family, and widespread diatoms *Synedra*, *Fragilaria*, *Navicula*, *Nitzschia* and green desmidian, protococcal algae.

Conclusions

1. In the studied water bodies (Tudakul lakes and Kuyumazar reservoir) of the Bukhara region, the following types of algae are more often found: *Bacillariophyta* (34 species); *Chlorophyta* (20 species); *Cyanophyta* (15 species); less often - *Dinophyta* (5 species); *Englenophyta* (1 species).
2. It was revealed that the highest total phytoplankton abundance in water samples from both reservoirs was *Cyanophyta* (6500.00×10^3 cells / l and 706.250×10^3 cells / l, respectively), and the highest phytoplankton biomass was noted in *Bacillariophyta* (187.800 mg / l) and *Chlorophyta* (188.400 mg / l). At the same time, *Englenophyta* and *Dinophyta* were not found in water samples from the Kuyumazar reservoir.
3. It was proved that in the phytoplankton samples from Lake Tudakul, planktonic freshwater-brackish-water forms b-mesosaprobic predominated; brackish-water b- and b- and α -mesosaprobic species of algae prevailed.

LIST OF USED LITERATURE

1. Жумаева, Ш. Б. (2022). КОЛИЧЕСТВЕННЫЙ УЧЕТ И КАЧЕСТВЕННАЯ ХАРАКТЕРИСТИКА ФИТОПЛАНКТОНА В ВОДОЕМАХ БУХАРСКОЙ ОБЛАСТИ. *Scientific progress*, 3(1), 1132-1136.
2. Худайкулова, Н. И., & Жумаева, Ш. Б. (2020). О стимуляции иммунитета на гиалуронидазу-фактор патогенности паразитов. In *Университетская наука: взгляд в будущее* (pp. 106-108).
3. Mustafaeva, M. I., & Jumayeva, S. B. (2020). Ecofloristic analysis of natural algae population of reservoirs used as biological ponds of Bukhara. In *Университетская наука: взгляд в будущее* (pp. 260-266).
4. Sh, J., & Mustafaeva, M. I. (2018). Comparison of the algoflora of bioprides of bukhara with the analogue flora of the pond of uzbekistan. *AGRICULTURAL SCIENCES*, 12.
5. JUMAeva, S. <http://ijpsss.iscience.uz/index.php/ijpsss> Vol 2, Issue 2 2022.
6. Jumaeva, S. B., & Mustafaeva, M. I. (2020). HEALING PROPERTIES OF REPA. *Новый день в медицине*, (4), 514-515.
7. Nuraliev, N. A., Jumaeva, S. B., & Ginatullina, E. N. (2020). Basic qualitative and quantitative characteristics of phytoplankton in some water bodies of the Bukhara Region. *European Journal of Molecular and Clinical Medicine*, 7(7), 559-566.
8. ASLONOVA, M. (2022). DETERMINATION OF SUICIDALITY AGAINST THE BACKGROUND OF PARASITIC DISEASES IN CHILDREN. *International Journal of Philosophical Studies and Social Sciences*, 2(2), 9-12.
9. Ramazonovna, A. M. (2022). CASES OF HYPOVITAMINOSIS ON THE BACKGROUND OF HELMINTHIASIS. *Web of Scientist: International Scientific Research Journal*, 3(02), 305-312.
10. Raximovich, O. A., & Ramazonovna, A. M. (2022). TO IDENTIFY THE CALCIOLOGICAL DEFICIENCY THAT OCCURS AGAINST THE BACKGROUND OF HELMINTHIASIS IN CHILDREN. *CENTRAL ASIAN JOURNAL OF MEDICAL AND NATURAL SCIENCES*, 3(2), 91-95.
11. Кароматов, И. Д., & Аслонова, М. Р. (2018). Лечебные свойства растения мак самосейка (обзор литературы). *Биология и интегративная медицина*, (2), 101-110.
12. Кароматов, И. Д., & Аслонова, М. Р. (2018). Малина как лечебное средство. *Биология и интегративная медицина*, (3), 211-226.
13. Nazarov, A. I. (2022). CONSEQUENCES OF SEIZURES AND EPILEPSY IN CHILDREN. *Web of Scientist: International Scientific Research Journal*, 3(02), 483-489.
14. Nazarov, A. I. (2022). ATROF-MUHITNING INSON SALOMATLIGIGA TA'SIRI. *Scientific progress*, 3(1), 881-885.
15. Ilhomovich, N. A. (2022). CLINICAL AND EPIDEMIOLOGICAL FEATURES OF COVID-19 DISEASE IN CHILDREN. *BARQARORLIK VA YETAKCHI TADQIQOTLAR ONLAYN ILMIY JURNALI*, 2(9), 182-186.
16. Ilhomovna, N. F. (2022). Negative Impact of Seizures on Quality of Life. *Miasto Przyszłości*, 24, 120-122.
17. Ilhomovna, F. N. (2022). LATE SEIZURES AND CONSEQUENCES OF EPILEPSY IN YOUNG CHILDREN. *Conferencea*, 219-223.
18. Ilhomovna, F. N. (2022). RESPONSIBILITY OF PARENTS BEFORE THE OFFSPRING. *Conferencea*, 441-446.
19. Rustamovna, T. N. (2020, December). COMPARATIVE FEATURES OF MORPHOMETRIC CHANGES IN THE ULTRASONIC ANATOMY OF THE THYROID AND PHYSICAL

- DEVELOPMENT OF CHILDREN 9 YEARS OF AGE. In *Archive of Conferences* (Vol. 12, No. 1, pp. 5-6).
20. Temirova, N. R. (2021, January). COMPARATIVE FEATURES OF ULTRASOUND ANATOMY OF THE THYROID GLAND AND ANTHROPOMETRIC PARAMETERS IN CHILDREN 7 YEARS OF AGE. In *Archive of Conferences* (pp. 32-33).
21. Rustamovna, T. N. (2021). Ultrasound anatomy of the thyroid gland children 4–5 years old living in the bukhara region. *ACADEMICIA: AN INTERNATIONAL MULTIDISCIPLINARY RESEARCH JOURNAL*, 11(2), 348-351.
22. Nazarova, D. (2019). POETRY OF JAMAL KAMOL. *Scientific Bulletin of Namangan State University*, 1(7), 186-189.
23. Назарова, Д., & Зойирова, Д. А. (2022). ЖАМОЛ КАМОЛ ДОСТОНЛАРИНИНГ БАДИИЙ ҚУРИЛИШИ. *Scientific progress*, 3(1), 934-944.
24. Nazarova, D. I. (2020). Jamol Kamolning Asru radifli g'azaliga yozgan muxammasi. *Ilim ham ja'miyet. Ilmiy-uslubiy jurnal*, (1), 93.
25. Akmalovna, A. C. (2022). Characteristics and Advantages of Soybean Benefits in Every way. *Journal of Ethics and Diversity in International Communication*, 1(8), 67-69.
26. Aminjonova, C. A. (2022). SOG'LOM ONA VA BOLA–BAXTLI KELAJAK ASOSI. *Scientific progress*, 3(1), 874-880.
27. Худайкулова, Н. И., & Жумаева, Ш. Б. (2020). О стимуляции иммунитета на гиалуронидазу-фактор патогенности паразитов. In *Университетская наука: взгляд в будущее* (pp. 106-108).
28. Худойкулова, Н. И. (2018). Пути воспитания толерантности у молодежи. *Наука, техника и образование*, (11 (52)), 98-100.