



## Features of the Cerebral Hemodynamics, Microhemocirculation in Patients with Post-Covid Syndrome

R. M. Urinov, S. S. Pulatov

**Abstract:** Post-covid syndrome is a long-lasting (for 3 or more months) pathological manifestations after an acute period of the disease. The main signs include severe weakness, heaviness in the chest, feeling of incomplete inspiration, headaches, joint and muscle pain, sleep disturbance, depression, cognitive decline, thermoregulation disorder, etc. The mechanism of the post-covid syndrome may be associated with the emerging chronic inflammation of the vessels, which negatively affects the work of the nervous system in the first place, and also disrupts the functioning of internal organs. SARS-CoV-2 destroys the inner surface of blood vessels, which increases the risk of developing microthrombi in the microvasculature. COVID-19 can also have a direct destructive effect on the cells of the body, as well as cause an excessive immune response and provoke autoimmune diseases.

**Keywords:** cerebral, COVID-19, hemodynamics, syndrome

The aim of the study was to study the features of cerebral hemodynamics and microcirculation in patients with post-COVID syndrome.

**Materials and research methods.** The study included 100 young and middle-aged people from 30 to 55 years old, with a verified diagnosis of the consequences of the COVID-19 coronavirus infection, confirmed by laboratory methods of research and after negative results (PCR, ELISA) for SARS-CoV-2, after 15 - 35 weeks from the onset of the disease, which made up the main group. The patients of the main group were divided into subgroup I - 70 people who had a mild COVID-19 disease and subgroup II - 30 patients who had a moderate and severe disease complicated by pneumonia. The control group consisted of 20 healthy subjects of the appropriate sex and age composition who had not had a coronavirus infection or other viral infection over the past 6 months and had no acute and decompensated chronic pathology at the time of observation.

### Results of the study and their discussion.

The study of cerebral blood flow in patients with post-covid syndrome at rest revealed statistically significant differences in Vmax (maximum systolic velocity) blood flow in the main extracranial vessels of the carotid artery system and intracranial vessels of the carotid and vertebrobasilar systems.

In the background study of cerebral hemodynamic in patients of subgroups 1 and 2, multidirectional changes in the linear blood flow velocity (BFR) at the extracranial level were observed.

The rate of blood flow at rest in the common carotid artery (CCA) was significantly lower in patients of the first subgroup and was combined with venous dysgenic compared to the control group ( $p <$

0.001). Thus, the blood flow in the right CCA in patients of subgroup 1 was  $86.95 \pm 1.95$  cm/s and  $91.1 \pm 3.16$  in the control group ( $p < 0.001$ ); on the left CCA, the blood flow velocity in subgroup 1 was  $86.8 \pm 1.96$  cm/s and  $92.53 \pm 3.35$  cm/s in the control group ( $p < 0.001$ );

In patients of the second subgroup, there was an increase in speed indicators according to OSA -  $95.4 \pm 1.54$  cm/s, significantly relative to the indicator of the control group ( $p < 0.001$ ); on the left CCA, the blood flow velocity in the second subgroup was  $95.1 \pm 1.81$  cm/s ( $p < 0.01$ ), with the control group.

Indicators of the linear velocity of blood flow in the internal carotid artery (ICA) had similar trends with LBF in the CCA. The blood flow velocity was significantly lower in patients of the 1st subgroup compared to the control group ( $p < 0.001$ ). Thus, the blood flow in the right ICA in patients of subgroup 1 was  $52.58 \pm 2.07$  cm/s and  $62.6 \pm 4.57$  in the control group ( $p < 0.001$ ); in the left ICA, the blood flow velocity in subgroup 1 was  $52.75 \pm 2.2$  cm/s, and  $63.6 \pm 4.1$  cm/s in the control group ( $p < 0.001$ ); In patients of the second subgroup, there was an increase in speed indicators for ICA -  $65.2 \pm 1.73$  cm/s ( $p < 0.01$ ) relative to the control group; on the left LSA, the blood flow velocity in subgroup 2 was  $65.77 \pm 1.28$  cm/s, without significant differences with the control group.

When assessing the blood flow velocity in the vertebral artery (VA), it was also significantly lower in patients from subgroup 1, relative to the control group. Thus, the blood flow in the right VA in patients of subgroup 1 was  $12.72 \pm 0.84$  cm/s ( $p < 0.001$ ) with the control group  $13.93 \pm 1.53$ ; on the left VA, the blood flow velocity in subgroup 1 was  $12.74 \pm 0.84$  cm/s and  $14.40 \pm 1.28$  in the control group ( $p < 0.001$ ).

In patients with post-covid syndrome who had pneumonia in the acute period of the second subgroup, there was an increase in speed indicators on the right VA -  $13.17 \pm 1.06$  cm/s without significant differences with the control group; on the left VA, the blood flow velocity in subgroup 2 was  $13.39 \pm 1.18$  cm/s without significant differences from the control group.

The resistance indices (RI) were significantly reduced in the subgroup of patients of the first subgroup in OSA and PA and amounted to  $0.69 \pm 0.01$  and  $0.63 \pm 0.01$ , respectively ( $p < 0.01$ ). In patients from the second subgroup, an increase in RI in PA to  $0.67 \pm 0.01$  was observed in the 2A subgroup, which is significantly higher than the control group ( $p < 0.01$ ).

During the study, a violation of the anatomy of the vessels was noted in patients of the 2nd subgroup, with the early formation of a violation of the course of the vessels (C, S vessels), and the formation of initial signs of atherosclerotic vascular damage - a violation of differentiation into layers of the intima-media complex.

At the intracranial level, the linear velocity of blood flow in the middle cerebral artery (MCA) was assessed. A decrease in blood flow velocities was revealed both in the group of patients of the first and second subgroups of observation relative to the control group ( $p < 0.001$ ),

When assessing the venous outflow in the observation groups, all patients had signs of an increase in the velocity along the direct sinus to  $50 \pm 0.8$  and  $57 \pm 0.6$  cm/s, respectively, which were significant relative to the control group ( $p < 0.001$ ). In the veins of Rosenthal, the acceleration of shunting was up to  $27 \pm 0.98$  and  $32 \pm 0.55$  cm/s, respectively ( $p < 0.001$ ). The veins of the vertebral plexuses were dilated up to 5-6 mm, the shunt reached  $30 \pm 0.58$  cm/sec in both subgroups. In subgroup II, shunting through the middle cerebral vein was accelerated to  $20 \pm 0.78$  cm/sec, and in subgroup I, to  $17.3 \pm 0.4$  cm/sec ( $p < 0.001$ ).

In the control group, the discharge rates through the venous vessels remained within the age norms.

Thus, the study of cerebral hemodynamic in patients of the studied groups showed a significant effect of the venous component on autoregulation of cerebral blood flow. Thus, in patients who had a mild form of coronavirus infection, the velocity indicators of the venous link had significantly high values with reduced velocity indicators of the arterial vessels. Thus, a decrease in the velocity of linear blood flow through the main cerebral arteries, a decrease in the tone of arterial vessels was revealed, which is a compensatory reaction in case of difficulty in venous outflow, however, under

these conditions, perfusion cerebral insufficiency is possible, which leads to transient hypoxia of the brain and, as a result, impaired blood flow autoregulation.

## Literature

1. Antonovich Zh.V. Prolonged COVID: definition, mechanisms, clinical manifestations, approaches to diagnosis and treatment. Focus on cough / Prescription. - 2021. - Volume 24 No. 4, part 2. S. - 23-44.
2. Temporary guidelines. Medical rehabilitation in case of a new coronavirus infection (COVID-19), M3 RF, Version 3 / Physical and rehabilitation medicine, medical rehabilitation. - 2021. - T.Z, No. 1, Appendix 1.
3. The technique of a two-component model of neurovegetative and metabolic stabilization of patients with a complicated course of coronavirus infection COVID-19: a manual for doctors / A.N. Kondratiev, Yu.S. Aleksandrovich, N.V. Dryagina, N.A. Lesteva, D.M. Rizakhanov, L.M. Tsenciper. - St. Petersburg: Association of Anesthesiologists-Resuscitators of the North-West, 2020. - 24 p.
4. Pulatov S.S., Urinov R.M. General clinical and neurological assessment in patients with post-COVID syndrome. Journal of neurology and neurosurgery research / VOLUME 3, ISSUE 5. 2023, 81-84.
5. Correction of mitochondrial dysfunction in the complex rehabilitation of patients after COVID-19 / Tereshin A.E. and others. II Journal of Neurology and Psychiatry named after S.S. Korsakov. - 2021. -t. 121, No. 8.-S. 25-29.
6. Chan, Y. The role of acupuncture during the COVID-19 pandemic: From the historical perspective to practical application / Y.Chan // Traditional Medicine and Modern Medicine Vol. 3, no. 1 (2020) 27-35.
7. Luks, Andrew & Swenson, Erik. (2020). Pulse Oximetry for Monitoring Patients with COVID-19 at Home: Potential Pitfalls and Practical Guidance. Annals of the American Thoracic Society. 17.10.1513/AnnalsATS.202005-418FR.
8. Pulatov SS, O'rinov RM Current Views on Neurological Disorders in Patients with Post-Cochlear Syndrome (Literature Review), International Journal of Health Systems and Medical Sciences. Volume 1 | #4 | Oct-2022 266-271
9. Ye, L. Analysis and Treatment of Early Symptoms of COVID-19 / L. Ye, P. Sun, T. Wang // EJOM Vol.9. No.5. 2020 TCM
10. Sanoeva M. et al. Comparative analysis of cognitive function at vascular complications of migraine (diagnosis and clinical approach) //Journal of Critical Reviews. – 2020. – Т. 7. – №. 3. – С. 425-430.
11. Саноева М. Ж., Саидвалиев Ф. С. Мигрень-вчера, сегодня, завтра. Современный взгляд на проблему //Международный неврологический журнал. – 2016. – №. 8 (86). – С. 72-78.
12. Sanoeva M. et al. Peculiarities of clinical and hemodynamic manifestations of migraine strokes //International Journal of Psychosocial Rehabilitation. – 2020. – Т. 24. – №. 2. – С. 350-358.
13. Саноева М. Ж., Жураева Г. Б., Мухидова Г. Х. Клинические особенности развития депрессии как предиктора осложненных форм мигрени //Вестник Совета молодых учёных и специалистов Челябинской области. – 2018. – Т. 1. – №. 3 (22). – С. 29-36.
14. Саноева М. Ж., Саидвалиев Ф. С., Гулова М. А. Современный взгляд к проблеме мигрени (обзорная статья) //Вестник Совета молодых учёных и специалистов Челябинской области. – 2016. – Т. 3. – №. 3 (14). – С. 59-66.
15. Саноева М. Ж., Саидвалиев Ф. С., Гулова М. А. Современный взгляд к проблеме мигрени (обзорная статья) //Вестник Совета молодых учёных и специалистов Челябинской области. – 2016. – Т. 3. – №. 3 (14). – С. 59-66.
16. Sanoeva M. Z., Rakhmatova D. I. Electromyographic Correlates and Non-Traditional Methods of Treatment of Complicated Forms of Facial Nervopathy //Биомедицина Ва Амалиёт Журнали. – С. 666.