



Effects of Sodium Glucose Cotransporters (SGLT2) Inhibitorstype 2 on Hepatosis in Patients Type 2 Diabetes Mellitus

Ismoilova Shaxnoza Samandarovna ¹

¹ Assistance of Bukhara State Medical Institution

Abstract: Empagliflozin is an antidiabetic agent used in adult patients with type 2 diabetes mellitus. It was FDA-approved in 2014. Empagliflozin can be used as a single agent or as a combination agent with other antidiabetic products. Combination products include empagliflozin and linagliptin and empagliflozin in combination with metformin. These newer agents can be more expensive for patients, thus impeding the clinician's ability to prescribe them based on patient financial considerations. However, the American Diabetes Association (ADA) is calling for agents with proven mortality reduction for use as second-line therapy after metformin. Such agents include empagliflozin and liraglutide. The primary treatment for diabetes is lifestyle management and exercise, followed by metformin use. Per Standards of Medical Care in Diabetes, if the A1c is greater than 9%, then combination therapy with metformin is recommended. In 2016, the FDA (United States Food and Drug Administration) approved a new indication for empagliflozin, which was to reduce the risk of cardiovascular death in adult patients with type 2 diabetes and cardiovascular disease. Empagliflozin has been shown to reduce hospitalizations for heart failure and death from cardiovascular causes. Patients are at an increased risk of cardiovascular mortality with type 2 diabetes, so prescribers should be made aware of the benefits of empagliflozin. In a nutshell, empagliflozin may be effective in the following settings.

Keywords: antidiabetic drug, clinical trial, liver, SGLT2 inhibitor.

Diabetes mellitus is a major public health disorder that affects more than 400 million subjects, with a prevalence ranging from 7.2 to 11.4% in the general population, worldwide [1, 2]. Diabetes is associated with various metabolic disorders and vascular damage of great extent [3] and thus, the management of the diabetic patient should aim in the attenuation of hyperglycemia and other concomitant cardiovascular conditions [4-7]. Current antidiabetic treatment options were unable to offer consistent cardiovascular benefits. Apart from metformin and glucagon-like peptide 1 analogues, the rest of the available hypoglycemic agents have not provided consistent benefits from the cardiovascular standpoint. Furthermore, the impact of these drugs on other cardiovascular risk factors is either neutral or beneficial, and in some cases even harmful [8-15]. Sodium-glucose cotransporters (SGLT) 2 inhibitors are a novel antidiabetic class of drugs that offer a different approach for the amelioration of hyperglycemia than the other antidiabetic drug classes. SGLT-2 inhibitors block the reabsorption of glucose in the renal proximal tubule, resulting in decreased glucose reabsorption, increased urine excretion of glucose and attenuation of glycemic parameters [6, 7]. Through their unique mechanism of action, SGLT-2 inhibitors seem to offer several beneficial

effects on a cluster of other cardiovascular risk factors. These drugs are associated with a mild increase in sodium urine excretion; this combined with the osmotic effect of the excreted glucose, results in a mild diuretic effect that leads to hemodynamic improvements. A mild decrease in blood pressure accompanied with a mild reduction in body weight is noted with SGLT-2 inhibition. In addition, beneficial effects with these drugs are observed in serum acid levels, triglycerides and high-density lipoprotein cholesterol levels. Ameliorating effects are also noted in other established cardiovascular risk factors such as diabetic nephropathy, non-alcoholic fatty liver disease and arterial stiffness [8, 9].

The pleiotropic effect of SGLT-2 inhibitors on the aforementioned abundance of cardiovascular risk factors has raised several expectations for their cardiovascular profile. As of this issue, two cardiovascular studies on these novel drugs exist and have provided encouraging results. Empagliflozin in the EMPA-REG study and (to a lesser extent) canagliflozin in CANVAS study have shown remarkable benefits in cardiovascular morbidity and mortality [10, 11]. Furthermore, SGLT-2 inhibitors seem to attenuate heart failure and renal outcomes [2, 3]. In this special issue of Cardiovascular and Hematological Diseases-Drug Targets, we aim to present current data on the multidimensional profile of SGLT-2 inhibitors and discuss the benefits and potential contradictory findings with this class of drugs. Manolis *et al.* discuss in detail the mechanism of action of SGLT-2 inhibitors and comprehensively explain the SGLT-2 inhibitors-induced increased excretion of glucose, sodium, water and of other substances that might result in important outcome benefits. They further describe the effect of SGLT-2 inhibition on other organs and systems, as well [2]. The antidiabetic profile of the drugs is presented by Kihm *et al.* in the following review. The authors report the results from studies of several SGLT-2 inhibitors used as monotherapy or as add-on treatment to preexisting antidiabetic drugs. They noted a significant reduction in fasting plasma glucose and glycated hemoglobin of approximately 0.5 to 1% (in absolute values). Importantly, it is stated that SGLT-2 inhibitors are not inferior to other hypoglycemic drugs in terms of glycemic control, with insignificant differences in glycated hemoglobin between SGLT-2 inhibitors-treated patients and other actively treated groups [5]. Subsequently, Tsioufis *et al.* in their review present the blood pressure-lowering effect of SGLT-2 inhibitors (mostly due to increased diuresis) from studies with office and ambulatory blood pressure measurements that range from 3 to 5 mmHg in systolic and 1 to 2 mmHg of diastolic blood pressure. Furthermore, it is stated that with their use, a mild body weight reduction of 2 to 3 kg is achieved, that is sustained with long-term treatment [6]. The significant cardiovascular and mortality risk reductions are reported by Lovic *et al.* who discussed the data obtained from the two aforementioned cardiovascular trials and from one large observational study in detail. Among the favorable data presented, the authors report the concerning higher amputation risk observed with canagliflozin use, as well [3]. Following this article, Doumas *et al.* and Papademetriou *et al.* comprehensively present data and potential underlying mechanisms of the attenuating impact of SGLT-2 inhibitors on renal and heart failure outcomes, respectively [10, 11]. In the article by Doumas *et al.* it is stated that SGLT-2 inhibitors are associated with a significant delay in the progression of diabetic kidney disease, with empagliflozin and canagliflozin achieving a remarkable deceleration of estimated glomerular filtration rate decline (0.9 ml/min/year) and amelioration of albuminuria status, respectively [8]. Papademetriou *et al.* present the benefits of SGLT-2 inhibition in heart failure outcomes and support that these findings could potentially be attributed to the ameliorating effect of SGLT-2 inhibitors on blood pressure, visceral obesity, arterial stiffness, diastolic dysfunction and importantly to the SGLT-2 inhibitors-induced increase in ketones bioavailability [7]. Tziomalos *et al.* report in detail the impact of these drugs on stroke risk factors and the contradictory neutral impact of SGLT-2 inhibitors on stroke risk, implying that other factors may be in play that need to be unveiled [3]. Last, Karagiannis *et al.* discuss the concerns of a potential increase in the risk of euglycemic diabetic ketoacidosis with this novel class of drugs. The authors present randomized controlled trials, meta-analyses, case series and case reports and note that most ketoacidosis cases were observed in patients with type 1 diabetes, due to off-label use of SGLT-2 inhibitors, or in patients previously misdiagnosed as having type 2 diabetes, who in fact suffered from late autoimmune diabetes of adults.

The recent CANVAS study provided similar results with the EMPA-REG trial and suggests a class effect of SGLT-2 inhibitors on morbidity and mortality outcomes. These remarkable findings could establish this class of drugs as an optimal second-line treatment option after metformin monotherapy.

The concerning findings in terms of stroke and amputations risk will place at the epicenter of the scientific interest, the identification of patients' groups that might be more prone to these events, such as patients with increased hematocrit levels or severe peripheral artery disease. Ongoing cardiovascular and renal trials will potentially provide further credence for this novel and very promising class of drugs. In this special issue of Cardiovascular and Hematological Diseases-Drug Targets, we aim to present current data on the multidimensional profile of SGLT-2 inhibitors and discuss the benefits and potential contradictory findings with this class of drugs. Manolis *et al.* discuss in detail the mechanism of action of SGLT-2 inhibitors and comprehensively explain the SGLT-2 inhibitors-induced increased excretion of glucose, sodium, water and of other substances that might result in important outcome benefits. They further describe the effect of SGLT-2 inhibition on other organs and systems, as well [24]. The antidiabetic profile of the drugs is presented by Kihm *et al.* in the following review. The authors report the results from studies of several SGLT-2 inhibitors used as monotherapy or as add-on treatment to preexisting antidiabetic drugs. They noted a significant reduction in fasting plasma glucose and glycated hemoglobin of approximately 0.5 to 1% (in absolute values). Importantly, it is stated that SGLT-2 inhibitors are not inferior to other hypoglycemic drugs in terms of glycemic control, with insignificant differences in glycated hemoglobin between SGLT-2 inhibitors-treated patients and other actively treated groups [5]. Subsequently, Tsioufis *et al.* in their review present the blood pressure-lowering effect of SGLT-2 inhibitors (mostly due to increased diuresis) from studies with office and ambulatory blood pressure measurements that range from 3 to 5 mmHg in systolic and 1 to 2 mmHg of diastolic blood pressure. Furthermore, it is stated that with their use, a mild body weight reduction of 2 to 3 kg is achieved, that is sustained with long-term treatment [11].

1 Type 2 diabetes further accelerates NAFLD progression from simple steatosis to non-alcoholic steatohepatitis (NASH), fibrosis and cirrhosis.² Moreover, NAFLD associates with an increased risk of cardiovascular diseases, including cardiomyopathy and certain cardiac arrhythmias, thereby contributing to the excess morbidity and mortality in people with both type 2 diabetes and NAFLD.² Hepatic fibrosis is a strong predictor of NAFLD-related mortality² and clinically relevant advanced fibrosis stages, F3 and F4, can be present in up to 20% of persons with NAFLD and type 2 diabetes.¹ While the 'gold-standard' diagnosis of fibrosis still requires liver biopsy,² several imaging methods have been introduced including vibration controlled transient elastography or magnetic resonance elastography. These methods are mostly restricted to use in specialized centres³ as they require specific technical equipment, qualified and trained personnel, making them difficult to perform in large multinational clinical trials involving hundreds of study sites. Thus, non-invasive indices calculated from demographic, anthropometric and laboratory parameters provide an opportunity to estimate the prevalence and effects of interventions on liver fibrosis in large cohorts.³ In addition, hepatic steatosis, the primary criterion of NAFLD diagnosis, may be estimated by non-invasive indices.^{4,5} Sodium glucose co-transporter 2 inhibitors are associated with a modest weight loss and improved cardiovascular and renal outcomes.^{6,7} Recent small-scale randomized controlled trials have shown that empagliflozin also improved hepatic steatosis and beneficial effects of empagliflozin on histological components including fibrosis were suggested from an uncontrolled pilot trial.⁶ The underlying mechanisms remain largely unknown; however, apart from weight loss, they may even include improvement of adipose tissue function with amelioration of local inflammation and/or oxidative stress.¹¹ The present study examined the effects of empagliflozin treatment in a large cohort of persons with type 2 diabetes with established cardiovascular disease (a) on indices of hepatic steatosis and fibrosis, (b) on glycaemia and body weight, as well as (c) on cardiorenal outcomes and all-cause mortality in groups at different steatosis and fibrosis risk.

Finally, this analysis addressed the question whether baseline steatosis and fibrosis risk scores are associated with the incidence of cardiorenal events in this patient population. To this end, an exploratory post-hoc analysis was performed in the EMPA-REG OUTCOME study, previously

showing lower rates of cardiovascular events and deaths from any cause with empagliflozin at a median observation time of 3.1 years.⁷ The design of EMPA-REG OUTCOME has been previously described.⁷ Adult individuals with type 2 diabetes and established cardiovascular disease were included; elevated liver enzymes >3x upper limit of normal were exclusion criteria. The main objective of this post-hoc analysis was to compare the effects of empagliflozin and placebo on Dallas steatosis index (DSI)¹⁰ and hepatic steatosis index (HSI)¹¹ as well as on NAFLD fibrosis score (NFS) and Fibrosis-4 score (FIB-4). All randomized participants, who received ≥ 1 dose of the study drug. The effects on risk scores as well as glycated haemoglobin (HbA1c) and weight were evaluated using a mixed-effect model repeated measurement model, which included baseline HbA1c and baseline of score (or weight) as linear covariates and their interaction with visit in addition to baseline estimated glomerular filtration rate (Modification of Diet in Renal Disease Study) category, geographical region and baseline body mass index category. Treatment, subgroup (if applicable) and visit were also entered as fixed effects as well as all two- and three-way interactions thereof. In addition, the model included a fixed categorical effect for ‘time of randomization’ to account for each patient’s theoretical ability to ‘reach’ certain weeks in this study arising from the study design. Because of small group sizes, participants at baseline intermediate and high risk (based on FIB-4) were pooled for analysis of time courses of parameters in FIB-4 low and high fibrosis risk categories and HSI low and intermediate risk categories were combined for all subsequent analyses. All time to first event analyses were performed with multivariate Cox regression models that included terms for sex, baseline age, estimated glomerular filtration rate, body mass index, HbA1c, geographical region, subgroup, and treatment _ subgroup interaction. Continuous baseline characteristics of fibrosis and steatosis risk groups are given as mean \pm standard deviation, categorical variables as number and proportions. All other data are expressed as adjusted means (95% confidence interval) or adjusted means \pm standard error.

The recent CANVAS study provided similar results with the EMPA-REG trial and suggests a class effect of SGLT-2 inhibitors on morbidity and mortality outcomes. These remarkable findings could establish this class of drugs as an optimal second-line treatment option after metformin monotherapy. The concerning findings in terms of stroke and amputations risk will place at the epicenter of the scientific interest, the identification of patients’ groups that might be more prone to these events, such as patients with increased hematocrit levels or severe peripheral artery disease. Ongoing cardiovascular and renal trials will potentially provide further credence for this novel and very promising class of drugs.

Conclusions: Empagliflozin may reduce steatosis but not fibrosis risk in individuals with type 2 diabetes and cardiovascular disease.

Literature:

1. International Diabetes Federation. *IDF Diabetes Atlas*, 7 ed.; International Diabetes Federation: Brussels, 2015.
2. Исмоилова Шахноза Самандаровна. (2023). Эффективность Применение Аналогов Инсулина При Сахарном Диабете 2 Типа С Кардиоваскулярными Осложнениями. Сравнительный Анализ Пациентов. *Central Asian Journal of Medical and Natural Science*, 4(4), 132-138.
3. Самандаровна, И. Ш. . (2023). Влияние Дефицита Витамина D На Метаболические Параметры У Женщин С Ожирением. *Research Journal of Trauma and Disability Studies*, 2(7), 51–60.
4. Бадридинова Б. RISK FACTORS FOR THE DEVELOPMENT OF DIABETIC NEPHROPATHY IN PATIENTS WITH TYPE 1 DIABETE //Журнал вестник врача. – 2020. – Т. 1. – №. 3. – С. 41-45.
5. САХАРНЫЙ ДИАБЕТ 2-ГО ТИПА У ДЕТЕЙ И ПОДРОСТКОВ БУХАРСКОЙ ОБЛАСТИ ХИЛОЛА ШАРИФОВНА ЯХЪЯЕВА
6. Биология и интегративная медицина, 139-145, 2021

7. Исмоилова Шахноза Самандаровна. (2023). Эффективность Применение Аналогов Инсулина При Сахарном Диабете 2 Типа С Кардиоваскулярными Осложнениями. Сравнительный Анализ Пациентов. *Central Asian Journal of Medical and Natural Science*, 4(4), 132-138.
8. XIII Яхёева ТА'ЛИМ ВА RIVOJLANISH TANLILI ONLAYN ILMİY JURNALI 3 (5), 319-321, 2023
9. Прогноз И Профилактика Диабетической Нефропатии ДР Худоёрова *Research Journal of Trauma Disability Studies* 2 (4), 75-84,2023
10. Особенности обмена витамина Д у женщин репродуктивного возраста на фоне ожирения.
11. BOSHQARUV VA ETIKA QOIDALARI ONLAYN ILMİY JURNALI 3 (11), 29-32, 2023
12. ВЛИЯНИЕ ДЕФИЦИТА ВИТАМИНА Д НА СОСТОЯНИЕ ЗДОРОВЬЯ ЖЕНЩИНЫ И ЕЁ РЕПРОДУКТИВНУЮ ФУНКЦИЮ (литературный обзор).ТА'ЛИМ ВА RIVOJLANISH TANLILI ONLAYN ILMİY JURNALI 3 (2), 1-4, 2023.
13. Исмоилова Ш. С. Влияние Дефицита Витамина D На Метаболические Параметры У Женщин С Ожирением //*Research Journal of Trauma and Disability Studies*. – 2023. – Т. 2. – №. 7. – С. 51-60.
14. Исмоилова, Шахноза Самандаровна. "Эффективность Применение Аналогов Инсулина При Сахарном Диабете 2 Типа С Кардиоваскулярными Осложнениями. Сравнительный Анализ Пациентов." *Central Asian Journal of Medical and Natural Science* 4.4 (2023): 132-138.
15. Ikhomovich A. N. Semirish Va Gipoterioz Kasalliklarining O'zaro Bog'liqlik Sabablari //AMALIY VA TIBBIYOT FANLARI ILMİY JURNALI. – 2023. – Т. 2. – №. 12. – С. 101-106.
16. Ikhomovich A. N. Pancreas in a Patient with Diabetes Mellitus the Current State of the Issue //American Journal of Pediatric Medicine and Health Sciences (2993-2149). – 2023. – Т. 1. – №. 9. – С. 501-506.
17. Rakhmatova D. B., Rakhmatova D. B. " Main" Symptoms and leading clinical options for the flow of acute coronary syndromes in women //Asian Journal of Multidimensional Research (AJMR). – 2019. – Т. 8. – №. 11. – С. 69-74.
18. Rakhmatova D. B. Analysis of the risk factors of Chd in persons over 60 years among the population of the city of Bukhara //Asian studies. Индия. – 2019. – Т. 1. – С. 33-38.
19. Rakhmatova D. B., Zikrillaev F. A. DETERMINE THE VALUE OF RISK FACTORS FOR MYOCARDIAL INFARCTION //FAN, TA'ЛИМ, MADANIYAT VA INNOVATSIYA. – 2022. – Т. 1. – №. 4. – С. 23-28.
20. Рахматова Д. Б., Аслонова М. Р. Современные методы медицинской реабилитации пациентов с артериальной гипертензией //Биология и интегративная медицина. – 2018. – №. 3. – С. 110-117.
21. Рахматова Д. Б. и др. ОЦЕНКА ФАКТОРОВ РИСКА ИБС У ЛИЦ СТАРШЕ 60 ЛЕТ СРЕДИ НАСЕЛЕНИЯ ГОРОДА БУХАРЫ //Теория и практика современной науки. – 2018. – №. 5 (35). – С. 704-708.
22. Рахматова Д. Б., Раджабова Г. Х. Анализ факторов риска ИБС у лиц старше 60 лет среди населения города Бухары //Биология и интегративная медицина. – 2019. – №. 3 (31). – С. 37-42.
23. Рахматова Д. Б. Гранат как лечебное средство в народной и древней медицине //Биология и интегративная медицина. – 2022. – №. 1 (54). – С. 157-168.

24. Rakhmatova D. B., Rakhmatova D. B. " Main" Symptoms and leading clinical options for the flow of acute coronary syndromes in women //Asian Journal of Multidimensional Research (AJMR). – 2019. – Т. 8. – №. 11. – С. 69-74.
25. Rakhmatova D. B. Scientific and practical significance of acute myocardial infarction among the population of elderly and old age //Globalization, the State and the Individual. – 2022. – Т. 29. – №. 1. – С. 84-89.
26. Sulstonova N. A. THE PROBLEM OF ADDICTED MISSING OF PREGNANCY IN EARLY STAGES OF PREGNANCY //Oriental Journal of Academic and Multidisciplinary Research. – 2023. – Т. 1. – №. 1. – С. 94-101.
27. Sulstonova N. A. Dopplerometric Features of Blood Flow Changes in the Utero-Placental System in Women With Related Pregnancy Mission //Miasto Przyszłości. – 2023. – Т. 34. – С. 268-273.
28. Sulstonova N. A. Evaluation of Clinical and Instrumental Results of Patients with a Risk of Development of Recurrent Mission //Central Asian Journal of Medical and Natural Science. – 2023. – Т. 4. – №. 2. – С. 536-542.
29. Султонова Н. А., Негматуллаева М. Н. Значимость Применения Витамина И Минеральной Комплексной Терапии В Профилактике Невынашивания Беременности //Central Asian Journal of Medical and Natural Science. – 2021. – С. 388-392.
30. Тиллоева Ш. Ш., Давлатов С. С. Эффективность и переносимость локсидола в лечение ревматоидного артрита у пациентов старших возрастных групп //Central Asian Journal of Medical and Natural Science. – 2021. – С. 432-436.
31. Тиллоева Ш. Ш. и др. Estimation of the condition of the cardiorespiratory system of patients with the conciliation of bronchial asthma and arterial hypertension, effects of complex therapy //Новый день в медицине. – 2020. – №. 2. – С. 227-230.
32. Tillaeva S. S. et al. Currency and diagnostic criteria of rheumatoid arthritis in patients of senior age groups //Asian Journal of Multidimensional Research (AJMR). – 2018. – Т. 7. – №. 11. – С. 184-188.
33. Ikhomovna K. D. Morphological Features of Tumor in Different Treatment Options for Patients with Locally Advanced Breast Cancer //International Journal of Innovative Analyses and Emerging Technology. – 2021. – Т. 1. – №. 2. – С. 4-5.
34. Khodzhaeva D. I. Changes in the Vertebral Column and Thoracic Spinecells after Postponement of Mastoectomy //International Journal of Innovative Analyses and Emerging Technology. – 2021. – Т. 1. – №. 4. – С. 109-113.
35. Khodjayeva D. I. MORPHOLOGY OF IDIOPATHIC SCOLIOSIS BASED ON SEGMENT BY SEGMENT ASSESSMENT OF SPINAL COLUMN DEFORMITY //Scientific progress. – 2022. – Т. 3. – №. 1. – С. 208-215.
36. Ikhomovna K. D. Modern Look of Facial Skin Cancer //BARQARORLIK VA YETAKCHI TADQIQOTLAR ONLAYN ILMIIY JURNALI. – 2021. – Т. 1. – №. 1. – С. 85-89.
37. Idiyevna S. G. Discussion of results of personal studies in the use ofmil therapy in the treatment of trauma to the oral mucosA //European Journal of Molecular medicineVolume. – Т. 2.
38. Idiyevna S. G. THE EFFECTIVENESS OF THE USE OF MAGNETIC-INFRARED-LASER THERAPY IN TRAUMATIC INJURIES OF ORAL TISSUES IN PRESCHOOL CHILDREN //Academic leadership. ISSN. – Т. 15337812.
39. Sharipova G. I. Light and laser radiation in medicine //European journal of modern medicine and practice. – 2022. – Т. 2. – №. 1. – С. 36-41.
40. Idievna S. G. THE EFFECT OF DENTAL TREATMENT-PROFILACTICS ON THE CONDITION OF ORAL CAVITY ORGANS IN CHILDREN WITH TRAUMATIC

STOMATITIS //Tibbiyotdayangikun» scientific-abstract, cultural and educational journal.- Bukhara. – 2022. – Т. 5. – №. 43. – С. 103-106.

41. Idievna S. G. CHANGES IN THE CONTENT OF TRACE ELEMENTS IN THE SALIVA OF PATIENTS IN THE TREATMENT OF PATIENTS WITH TRAUMATIC STOMATITIS WITH FLAVONOID-BASED DRUGS //Journal of research in health science. – Т. 6. – С. 23-26.
42. Ikromovna I. F., Jumatovich J. U., Elmuradovich I. G. Influence of the harmful factors of manufacture of synthetic detergents and cleaners on the clinical-functional parameters of the oral cavities in the workers //European science review. – 2014. – №. 9-10. – С. 31-32.
43. Ибрагимова Ф. И., Жумаева А. А., Ражабова Д. Б. Влияние неблагоприятных факторов условий труда в производстве синтетических моющих и чистящих средств на состояние тканей пародонта у рабочих //Наука молодых–Eruditio Juvenium. – 2015. – №. 1. – С. 31-34.
44. Ikromovna I. F. Prevalence and character of the oral cavity mucosa in the workers of the manufacture of the synthetic detergents //European science review. – 2016. – №. 3-4. – С. 178-179.
45. Ибрагимова Ф. И., Замонова Г. Ш. Влияние вредных факторов производства на клинко-функциональные показатели полости рта рабочих //Символ науки. – 2016. – №. 8-1. – С. 181-182.
46. Ibragimova F. I., Idiev G. E. The state of health of workers in the production of synthetic detergents and cleaning products." Problems of Biology and Medicine //International Scientific Journal.-Samarkand. – 2012. – №. 1. – С. 68.
47. Ибрагимова Ф. И., Жуматов У. Ж. Поражения слизистой оболочки полости рта у работающих в производстве синтетических моющих и чистящих средств //Молодежный инновационный вестник. – 2016. – Т. 5. – №. 1. – С. 165-166.
48. Ибрагимова Ф. И., Замонова Г. Ш. ОЦЕНКА НЕКОТОРЫХ ФУНКЦИОНАЛЬНЫХ ПОКАЗАТЕЛЕЙ ПОЛОСТИ РТА У РАБОЧИХ ПРОИЗВОДСТВА СЫРЬЕВЫХ КОМПОНЕНТОВ ДЛЯ СИНТЕТИЧЕСКИХ МОЮЩИХ СРЕДСТВ //Молодежный инновационный вестник. – 2016. – Т. 5. – №. 1. – С. 163-165.
49. Ибрагимова Ф. И., Замонова Г. Ш. Влияние вредных факторов производства на клинко-функциональные показатели полости рта рабочих //Символ науки. – 2016. – №. 8-1. – С. 181-182.
50. Ибрагимова Ф. И., Идиев Г. Э. Действие гипохлорита натрия (ингредиента синтетических моющих средств) на активность окислительно-восстановительных ферментов, и её коррекция введением растительных препаратов в эксперименте //Проблемы биологии и медицины–Самарканд. – 2017. – Т. 4. – С. 98.
51. Urinbaevna Y. R. Features of Prediction of the Severity of Iron Deficiency in Helicobacter Pylori Infection //Scholastic: Journal of Natural and Medical Education. – 2023. – Т. 2. – №. 4. – С. 93-99.
52. Юлдашова Р. У. ЭПИДЕМИОЛОГИЧЕСКАЯ ХАРАКТЕРИСТИКА ЖЕЛЕЗОДЕФИЦИТНОЙ АНЕМИИ У ДЕТЕЙ И ПОДРОСТКОВ В РЕСПУБЛИКЕ УЗБЕКИСТАН ЗА 2007-2019 ГОДЫ //Новый день в медицине. – 2020. – №. 4. – С. 742-747.
53. Юлдашова Р. У., Жарылкасынова Г. Ж., Сафоев Б. Б. МОДЕРНИЗАЦИЯ КУРСА ДОВРАЧЕБНОЙ НЕОТЛОЖНОЙ ПОМОЩИ В БУХАРСКОМ ГОСУДАРСТВЕННОМ МЕДИЦИНСКОМ ИНСТИТУТЕ КАК ОДИН ИЗ УСОВЕРШЕНСТВОВАННЫХ МЕТОДОВ ОБУЧЕНИЯ (в рамках проекта ModeNEd) //Оптимизация высшего медицинского и фармацевтического О-62 образования: менеджмент качества и инновации:

- материалы IX внутривузовской научно-практической конференции.—Челя-бинск: Издательство Южно-Уральского государственного меди-цинского университета, 2018.— 153,[1] с. – 2018. – С. 150.
54. Юлдашова Р. У. и др. ИСПОЛЬЗОВАНИЕ СИСТЕМЫ ДИСТАНЦИОННОГО ОБУЧЕНИЯ MOODLE ПРИ ПОВЫШЕНИИ КВАЛИФИКАЦИИ ВРАЧЕЙ //Оптимизация высшего медицинского и фармацевтического об-О-62 разования: менеджмент качества и инновации: материалы VIII внутривузовской научно-практической конференции.—Челя-бинск: Издательство Южно-Уральского государственного меди-цинского университета, 2017.—136 с. – 2017. – С. 135.
55. Юлдашова Р. У., Халилова Ф. А., Тошева Х. Б. ОТНОШЕНИЕ СТУДЕНТОВ И ПРЕПОДАВАТЕЛЕЙ К ИСПОЛЬЗОВАНИЮ СОЦИАЛЬНЫХ СЕТЕЙ В ОБУЧЕНИИ //Педагогический профессионализм в образовании. – 2015. – С. 218-219.
56. Ziyolloevich A. M. Etiotropic Therapy of Viral Hepatitis //EUROPEAN JOURNAL OF INNOVATION IN NONFORMAL EDUCATION. – 2022. – Т. 2. – №. 11. – С. 52-56.
57. Abdulloev M. Z. EFFECTIVENESS OF ANTIVIRAL THERAPY IN VIRAL HEPATITIS //Journal of new century innovations. – 2022. – Т. 11. – №. 3. – С. 132-136.
58. Абдуллоев М. З., Облокулов А. Р. АНАЛИЗ РЕЗУЛЬТАТОВ ИССЛЕДОВАНИЯ ПО ОПРЕДЕЛЕНИЮ ЦИТОКИНОВ У БОЛЬНЫХ ХРОНИЧЕСКОГО ГЕПАТИТА С В ЗАВИСИМОСТИ ОТ НАЛИЧИЯ КРИГЛОБУЛИНЕМИИ //“ XALQ TAVOVATI VA ZAMONAVIY TIBBIYOT, YANGI YONDASHUVLAR VA DOLZARB TADQIQOTLAR”. – 2023. – Т. 6. – С. 8-9.
59. Abdulloev M. Z. EFFECTIVENESS OF ANTIVIRAL THERAPY IN VIRAL HEPATITIS //Journal of new century innovations. – 2022. – Т. 11. – №. 3. – С. 132-136.
60. Ziyodulloevich A. M. EFFICIENCY OF ETIOTROPIC TREATMENT IN CHRONIC VIRAL HEPATITIS //Galaxy International Interdisciplinary Research Journal. – 2023. – Т. 11. – №. 4. – С. 450-454.
61. Oblokulov R. A., Abdulloev M. Z. To Study the Effectiveness of Direct-Antiviral Drug in the Treatment of Chronic Viral Hepatitis C, Your Act //Web of Scholars: Multidimensional Research Journal. – 2022. – Т. 1. – №. 5. – С. 201-206.
62. Облокулов А. Р. и др. ВИРУСОЛОГИЧЕСКИЙ ОТВЕТ ПРИ ЛЕЧЕНИИ ВИЧ-ИНФЕКЦИИ АНТИВИРУСНЫМИ ПРЕПАРАТАМИ //Новый день в медицине. – 2020. – №. 1. – С. 306-308.
63. Ахмедова С. М. Морфологическая характеристика развития стенок сердца крысят //Наука и мир. – 2015. – №. 1-2. – С. 85-87.
64. Каримов Х., Ахмедова С., Тен С. Морфологическая характеристика развития стенок сердца и их изменения при воздействии пестицидов //Журнал вестник врача. – 2011. – Т. 1. – №. 03. – С. 51-54.
65. Ахмедова С., Нортюева Н., Нортюев А. Morphological changes in the teeth of adolescent children with hypotireosis : дис. – Тошкент тиббиёт академияси, 2022.
66. Ахмедова С. М. Возрастные особенности анатомии сердца крысят в раннем постнатальном онтогенезе //International medical scientific journal. – 2015. – С. 40.
67. Ахмедова С. М. Гистотопография стенок сердца крысы в постнатальном онтогенезе //Врач-аспирант. – 2011. – Т. 46. – №. 3.2. – С. 283-288.
68. Ахмедова С. М., Айтжанова А. Е., Сагдуллаева М. К. К МОРФОЛОГИИ ИЗМЕНЕНИЙ ПОЧЕК ПРИ ЭКСПЕРИМЕНТАЛЬНОМ АЛКОГОЛИЗМЕ //Journal of new century innovations. – 2022. – Т. 16. – №. 2. – С. 166-168.

69. Ахмедова С. М., Якубова Ф. ПОКАЗАТЕЛИ ФИЗИЧЕСКОГО РАЗВИТИЯ У СЛЕПЫХ И СЛАБОВИДЯЩИХ ДЕТЕЙ //Conferencea. – 2022. – С. 103-105.
70. Миршарапов У. М. и др. СОСТОЯНИЕ СОСУДОВ ПРИ ЭКСПЕРИМЕНТАЛЬНОМ САХАРНОМ ДИАБЕТЕ //Проблемы и достижения современной науки. – 2017. – №. 1. – С. 13-15.
71. Ахмедова С. М., Хатамов А. И., Худайбергенов Б. Э. МОРФОФУНКЦИОНАЛЬНЫЕ ОСОБЕННОСТИ УЛЬТРАСТРУКТУРНОГО СТРОЕНИЯ МИОКАРДА ПРИ ВЛИЯНИИ ПЕСТИЦИДА //Проблемы и достижения современной науки. – 2017. – №. 1. – С. 11-13.
72. Ахмедова С., Эрматов Н. Характер морфофункциональных изменений сердца при воздействии пестицидов //Журнал проблемы биологии и медицины. – 2016. – №. 3 (89). – С. 135-137.
73. Pkhomovich A. N. Semirish Va Gipoterioz Kasalliklarining O'zaro Bog'liqlik Sabablari //AMALIY VA TIBBIYOT FANLARI ILMIY JURNALI. – 2023. – Т. 2. – №. 12. – С. 101-106.
74. Pkhomovich A. N. Pancreas in a Patient with Diabetes Mellitus the Current State of the Issue //American Journal of Pediatric Medicine and Health Sciences (2993-2149). – 2023. – Т. 1. – №. 9. – С. 501-506.
75. Nasirova S. Z. Polypharmacy As An Actual Problem Of Pharmacotherapy //The American Journal of Medical Sciences and Pharmaceutical Research. – 2021. – Т. 3. – №. 01. – С. 1-5.
76. Nasirova S. Z. MORPHOMETRIC PARAMETERS OF THE LIMPHOID TISSUE OF THE SMALL INTESTINE WHEN USING ANTI-INFLAMMATORY DRUGS //Asian journal of pharmaceutical and biological research. – 2022. – Т. 11. – №. 2.
77. Zauravna N. S. EFFECTS AND ACTIONS OF SILYBUM MARIANUM PHYTOPREPARATION //Научный Фокус. – 2023. – Т. 1. – №. 3. – С. 300-308.
78. Nasirova S. Z. Effect of anti-inflammatory medicines on the morphometric structure of the peyer's patches on the small intestine //Modern views and research. International scientific and practical Conference Egham.-England. – 2021. – С. 85-86.
79. Nasirova S. Z., Samadov A. T. Changes in morphometric parameters of the small intestine in the conditions of polypragmasy //Тиббиётда янги кун. – 2021. – Т. 2. – №. 34/1. – С. 28-32.
80. Nasirova S. Z. Changes in morphometric parameters of the lymphoid tissue of the small intestine in the conditions of polypragmasia //American Journal of Medicine and Medical Sciences.- America N. – 2021. – Т. 11. – С. 673-677.
81. Zauravna N. S. MAIN EFFECTS OF SÍLYBUM MARIÁNUM //Asian journal of pharmaceutical and biological research. – 2023. – Т. 12. – №. 1.