



Growth and Hormonal Status of Children in the Context of Reforming the Preschool Education System

Yadgarova Shaxodat Solixovna ¹

¹ Assistant of the Department of Ecology and Hygiene of the Bukhara State Medical Institute

Abstract: Growth hormone (GH) deficiency is the most common pituitary hormone deficiency in children and can occur in isolation or in combination with other pituitary hormone deficiencies. In typical cases, growth hormone deficiency in children is characterized by an abnormally low growth rate and short stature while maintaining normal body proportions. Diagnostics include measuring pituitary hormone levels and CT or MRI to detect structural abnormalities of the pituitary gland or brain tumors. Treatment usually includes hormone replacement therapy and removal of the tumor if it is causing the disorder.

Keywords: Growth hormone, panhypopituitarism, GH-releasing hormone receptor, hereditary genetic causes.

There is both an isolated deficiency of growth hormone, and associated with pan hypopituitarism. In either case, GH deficiency can be acquired or congenital (including due to hereditary genetic causes). In rare cases, there is an abnormality of the GH receptors, which causes insensitivity to the hormone at normal levels in the blood.

Isolated GH deficiency occurs with a frequency of 1/4000 to 1/10000. As a rule, it is idiopathic, but about 25% of cases have an established etiology. Congenital causes include abnormalities of the GH-releasing hormone receptor and GH1 genes, as well as some malformations of the central nervous system. Acquired causes include therapeutic radiation to the central nervous system (high doses of radiation can cause pan hypopituitarism), meningitis, histiocytosis, and traumatic brain injury. Preventive or therapeutic irradiation of the spine further reduces the growth potential of the vertebrae and threatens to further reduce the final growth.

Pan hypopituitarism can have genetic causes, such as hereditary or sporadic mutations that affect cells in the pituitary gland. In such cases, pathologies of other organs and systems are also encountered; in particular, midline facial defects such as cleft palate or septooptic dysplasia (which includes septum atresia, optic atrophy and hypopituitarism). Panhypopituitarism can also result from a variety of lesions that affect the hypothalamus (disrupting the secretion of releasing hormones) or the pituitary gland, examples include tumors (eg, most commonly craniopharyngioma), infections (eg, tuberculosis, toxoplasmosis, meningitis), and infiltrative processes. The combination of lytic damage to the bones of the skeleton or skull with diabetes insipidus indicates Langerhans cell histiocytosis.

About 3% of children have severe growth retardation. At the same time, growth hormone deficiency as a cause of short stature is detected in no more than 8.5% of them. In other children, the most often revealed constitutional features of growth and development, less often - a deficiency of other anabolic hormones, severe somatic diseases, genetic and chromosomal diseases.

Stunting can be caused by two fundamentally different problems. The first is when the child is stunted due to an existing disease. Then treatment is needed, which, quite possibly, will really allow him to grow up and maintain health. Which of these factors affect growth? Infectious

diseases, heart defects, chronic bone diseases, etc. cause various disorders in the body and retard its growth.

Diseases of the endocrine glands, such as the pituitary gland, the thyroid gland, and the adrenal glands, have a particularly great influence. That is why growth retardation should be the reason for contacting an endocrinologist. The most severe disorders of growth processes are observed in pathology of the endocrine system. It is known that almost all hormones are directly or permissively involved in growth processes.

Human growth is genetically programmed. It is carried out by somatotropin. It is called growth hormone because in children and adolescents, as well as young people with not yet closed growth zones in the bones, it causes a pronounced acceleration of linear (in length) growth, mainly due to the growth of long tubular bones of the extremities. Growth hormone secretion gradually decreases with age. It is minimal in the elderly and the elderly, maximum in adolescents during the period of intense linear growth and puberty. In adults, a pathological increase in the level of growth hormone or prolonged administration of exogenous growth hormone at doses characteristic of a growing organism leads to thickening of the bones and coarsening of facial features. If the secretion of this hormone stops prematurely, then growth stops. This can be for various reasons: injury, diseases, etc. Growth retardation (nanism) caused by a deficiency of growth hormone is one of the urgent problems of the whole society. The disease manifests itself in a significant slowdown in physical development; as a result, children who do not receive appropriate treatment cannot reach the usual average height of an adult over the years and are doomed to remain small throughout their lives, in fact - disabled.

In most cases, growth retardation is due to hereditary factors. Children of short parents with the so-called family delayed maturation (puberty begins later than usual) grow poorly. Usually, this requires only the supervision of a doctor, but sometimes in these cases the help of an endocrinologist may be required: with age, such children, due to slower growth or late onset of puberty, may not reach their maximum possible growth potential inherent in the genes. Currently, with a timely visit to a doctor, this can be avoided and even slightly improved growth rates. In some families, there is a low growth in both parents, then the growth retardation in a child in the absence of signs of any pathology is just a constitutional feature. With such a short stature, no special treatment is required, however, general strengthening measures will help the body realize its existing potential. In the first years of life, growth retardation can occur due to malnutrition, impaired intestinal absorption. If the child is lagging behind in growth from peers, and the reason for the lag is unclear, be sure with the help of a doctor to exclude the latent course of celiac disease in the child - intolerance to gluten (gluten) - a protein found in wheat and some other cereals. Depending on the cause of growth retardation, they are divided: endocrine - associated with a disease of the endocrine glands (for example, underdevelopment or lack of development of the pituitary gland (area of the brain), hypothyroidism (decreased levels of thyroid hormones), etc.); non-endocrine - not associated with endocrine gland disease.

References

1. Meerson F.Z. On the "price" of adaptation. *Patol. physiology and experimental therapy*. 1986; 3: 9-19.
2. Olimova D. V. A complex approach to glossalgia treatment based on the current data on the specificity of its etiopathogenesis //integration of science, education and practice. *Scientific-methodical journal*. – 2022. – T. 3. – №. 2. – C. 141-146.
3. Fayzullayevich, S. S., Hikmatovna, K. S., Khasanovna, M. M., & Ul'yanovna, S. G. (2018). Immune status in patients with duodenal ulcer and influence on her immunomodulatory therapy. *European science review*
4. ОЛИМОВА, Д. В. "USE OF MODERN METHODS IN THE TREATMENT OF GLOSSALGIA." *Analytical Journal of Education and Development*, <https://sciencebox.uz/index.php/jars/issue/view/45> P.-197-200.

5. Salikhovna Y. S. Basics of Hygienic Assessment of Children's Physical Development in Preschool Educational Institutions //AMALIY VA TIBBIYOT FANLARI ILMIY JURNALI. – 2022. – C. 119-123.
6. Nazarov J., Sharipova M. A. The Immune System in Children with Different Incidence Rates //Central asian journal of medical and natural sciences. – 2021. – C. 5-7.
7. Nazarov J. S. E. et al. Diuretic Therapy of Ascites in Patients with Liver Cirrhosis in the Absence of Azotemia //central asian journal of medical and natural sciences. – 2021. – C. 22-24.
8. Salikhovna, Yadgarova Shakhodat. "scientific and methodological bases of a doctor's activity in hygiene of children and adolescents." *barqarorlik va yetakchi tadqiqotlar onlayn ilmiy jurnali* (2022): 132-135.
9. Nazarov Z. S. E. Using brainstorming and case-study method in practical classes of microbiology //Новый день в медицине. – 2021. – №. 1. – C.
10. Suleymanov, S. F., Mansurova, M. K., Sagdullayeva, G. U., & Nazarov, J. S. E. (2019). Immune status in patients with duodenal ulcer and influence on her immunomodulatory therapy. *Asian Journal of Multidimensional Research (AJMR)*, 8(12), 84-87.
11. Ortikov A. A. Some hygiene issues according to the conditions of the workers of poultry farms //ACADEMICIA: An International Multidisciplinary Research Journal. – 2021. – T. 11. – №. 3. – C. 496-501.
12. Akhrorovich O. A. Poultry farm as a source of environmental pollution //ACADEMICIA: An International Multidisciplinary Research Journal. – 2021. – T. 11. – №. 11. – C. 554-558.
13. Akhrorovich O. A. Hygienic assessment of working conditions and prevention of reproductive pathology among workers of the poultry complex //ACADEMICIA: An International Multidisciplinary Research Journal. – 2022. – T. 12. – №. 1. – C. 161-166.
14. Oxunjanova, M. Z., and Akhrorova Sh B. Features of neurosensory. "disorders in patients with diabetic polyneuropathy and the possibility of nondrug correction." *Web of Scientist: International Scientific Research Journal (WoS)* 3.1 (2022): 818-825.
15. Zafarovna O. M. Negative Effect of Industrial Dust on the Human Body //AMALIY VA TIBBIYOT FANLARI ILMIY JURNALI. – 2022. – C. 128-131.
16. Samandarovna S. K. The Problem of Environmental Pollution //Vital Annex: International Journal of Novel Research in Advanced Sciences. – 2022. – T. 1. – №. 3. – C. 81-85.
17. Samandarovna S. X. Mehnat Sharoitining Inson Organizmiga Ta'siri Sog 'Lom Turmush Tarzi //AMALIY VA TIBBIYOT FANLARI ILMIY JURNALI. – 2022. – C. 34-37.