



Study of the Specificity of the Immunological State in Kidney Transplantation

Ismoilov Ibodjon Imomjonovich ¹

¹ Bukhara State Medical Institute named after Abu Ali Ibn Sino

Abstract: During this period, tremendous experience has been accumulated regarding the modernization of surgical techniques, organ preservation, improvement and optimization of immunosuppression protocols, as well as postoperative management of patients. Compared to transplantation of other organs, such a direction as kidney transplantation has more than half a century of history. By the end of the 90s, modern survival rates for renal transplants and recipients had been achieved. The success of kidney transplantation, however, has led to the fact that "waiting lists" for the operation are growing steadily every year around the world.

Keywords: Immunosuppression, hemodialysis, allogeneic transplant, immune response.

Treatment of end-stage chronic renal failure is one of the most pressing modern medical and social problems. Existing methods of renal replacement therapy for end-stage chronic renal failure: hemodialysis. Peritoneal dialysis and allotransplants of the kidneys are constantly being improved, which leads to an increase in the life expectancy of patients, but not always - in quality. At the same time, the number of patients requiring renal replacement therapy is constantly increasing, which complicates the availability of J10. Today, kidney transplantation is considered as the optimal method of renal replacement therapy, since it increases the duration and quality of life of patients to a greater extent than peritoneal and hemodialysis. In addition, it is known that kidney transplantation provides a higher quality of life for patients, and is also the most preferred by the method of renal replacement therapy from an economic point of view [5],

One of the essential factors determining the success of organ transplantation is the immunological compatibility of donor and recipient tissues. Achievements of the last decade of the last century in the field of fundamental immunogenetics, which made it possible to concretize the mechanisms of the implementation of the main functions of proteins. The gene-encoded immune response, as well as the transition from the study of genes of the human immune response from serological to molecular-genetic, have not only opened up fundamentally new possibilities for introducing these advances into medicine, but have made it possible to put them into practice with unprecedented speed [4].

Thanks to these advances, the effectiveness of kidney transplants (based on the results of the annual survival rate in the centers that switched to donor selection based on molecular genetic typing) increased by twenty%. At the same time, the remaining indicators of the functional graft significantly improved.

The discovery and use of modern methods of immunosuppression made it possible to reduce the likelihood of developing an acute rejection reaction and to increase the graft survival rate during the first year after the operation, leaving the long-term survival rates of the renal allograft unsatisfactory. In connection with this, methods of predicting the conflict "donor-recipient" are being actively developed.

The interest of researchers in the study of immunological parameters in organ transplantation and kidney transplantation in particular has always remained at a high level, but the opportunity to study the delicate relationship between the recipient's organism and the donor organ has appeared in the last twenty years. This is associated with a number of important discoveries in the field of fundamental immunology. This entailed the introduction into wide practice of new reliable and highly sensitive research methods, modern instruments and diagnostic systems [2]. The use of new complex methods requires clinical understanding.

So, in particular, after the introduction into practice of the molecular genetic approach to typing HLA genes, it became necessary to conduct a comparative analysis of the results obtained by both methods. Such studies were carried out by many laboratories around the world that are engaged in histotyping [1]. The data obtained revealed serious discrepancies in the typing results. Therefore, similar works aimed, ultimately, at the optimization of immunophenotyping continue and remain relevant.

Treatment of chronic renal failure is one of the medical and social problems for modern medicine. There are now a growing number of methods for the kidney and its maintenance, for example: Temporary renal hemodialysis, in addition to kidney transplantation, which increases the life expectancy of patients. But not always. Currently, the number of patients in need of kidney transplantation and the number of transplant options is increasing.

Renal transplantation, renal complications, impaired adaptation of the donor organism to the transplanted kidney are associated with the immunogenesis of the individual organism. Changes in the physiological activity of class T lymphocytes in the body after kidney transplantation are accompanied by changes in all immunogenetic conditions in the body, which leads to a decrease in renal vital signs, resulting in renal complications, decreased vital signs of renal transplantation within three years and changes in physiological functions. . In this regard, we organize research, observational work on the preservation of vital signs of the kidney, placed in the state of studying its immunogenesis after kidney transplantation.

It is known that a graft transplanted to a recipient from a genetically foreign donor does not take root and is inevitably rejected. At the same time, genetic differences between donor and recipient tissues play a key role in the development of allogeneic transplant rejection.

Antigens providing intraspecific differences are designated as tissue compatibility (histocompatibility) antigens and belong to the major histocompatibility gene complex (MHC) [6]. In humans, the MHC is called HLA (human leukocyte antigen). The biological significance of MHC lies in ensuring the interaction of body cells, recognizing its own, foreign and altered own cells, triggering and implementing an immune response against carriers of foreign information, positive and negative selection of T-cell clones, presentation of the targets of the immune response.

The immunological nature of graft rejection was demonstrated by Peter Medawar in an experiment on the transplantation of a genetically alien skin graft in rabbits [7]. Both humoral and cellular mechanisms play a role in transplant rejection. Cellular rejection mechanisms cause T-lymphocytes to become sensitized to the transplanted antigens. These lymphocytes cause damage to cells of foreign tissue by either direct cytotoxicity or secretion of lymphokines. T cell damage is characterized by parenchymal cell necrosis, lymphocytic infiltration, and fibrosis. Humoral mechanisms are mediated by antibodies that may be present in the serum of the recipient before transplantation or develop after transplantation of foreign tissue. Humoral factors damage the transplanted tissue through reactions that are equivalent to type II and III hypersensitivity reactions. The interaction of antibodies with the antigen on the surface of the transplanted cells leads to cell necrosis, and the accumulation of immune complexes in the blood vessels activates complement, which leads to the development of acute necrotizing vasculitis or chronic fibrosis of the intima with vasoconstriction.

The tolerance of the immune system is understood as a specific immunological nonresponsiveness to antigens. In this case, the absence of a response to this antigen is characteristic, but the response to any other is preserved. According to the figurative expression of R.V. Petrova, tolerance is immunity

with a minus sign. The lack of response of the immune system to its own antigens protects the body from autoaggression [8]. When tolerance to alloantigens is established, the transplanted tissue may take root. Tolerance to antigens exogenously entering the body can be induced both during the neonatal period and at puberty. The mechanisms of the immune system that allow blocking aggression against one's own or donor cells and tissues are conditionally divided into central and peripheral. Central tolerance is induced in the central organs of immunogenesis - in the thymus gland and bone marrow - and limits the autoreactivity of T and B lymphocytes.

References

1. II Imomjonovich, SS Fayzullayevich, NJS Erkinovich. Immunogenesis of Kidney Transplantation, Maintenance of Vital Signs of Transplanted Kidney. *Annals of the Romanian Society for Cell Biology*. 2021/3/30.
2. Imomjonovich I. I., Amirkulovna A. G. Methods of early detection of rejection in a kidney transplant from a relative donor // *Academicia Globe: Inderscience Research*. – 2021. – Т. 2. – №. 05. – С. 293-295.
3. Raychaudhuri S.P., Kundu-Raychaudhuri S., Tamura K., et al. FR255734, a humanized, Fc-Silent, Anti-CD28 antibody, improves psoriasis in the SCID mouse-psoriasis xenograft model. *J. Invest. Dermatol.* 2008;128:1969-1976. PMID:18337836 DOI:10.1038/jid.2008.38
4. Hikmatovna N. Z. Optimization of treatment of early neurological complications in cardioembolic stroke // *Middle European Scientific Bulletin*. – 2021. – Т. 8.
5. Hikmatovna B. Z., Zhalilovna H. D. Modern principles of management of patients with lower back pain // *Biology and Integrative Medicine*. – 2020. – №. 2. – С. 42.
6. Hikmatovna B. Z., Tukhtaeva K. K. Abu ali ibn sina about a healthy lifestyle // *Academicia Globe: Inderscience Research*. – 2021. – Т. 2. – №. 6. – С. 380-383.
7. Babajanova Z. K. The quality of clinical examination of patients with neurological diseases in family polyclinics // *Asian journal of multidimensional research*. – 2021. – Т. 10. – №. 4. – С. 1008-1012.
8. Le Blanc K. Immunomodulatory effects of fetal and adult mesenchymal stem cells. *Cytherapy*. 2003; 5(6) :48 5 — 48 9. PMID: 146 60044 DOI:10.1080/14653240310003611
9. Isroilovich A. E. et al. The Role And Importance Of Glioh Neurotrophical Factors In Early Diagnosis Of Parkinson Disease // *Texas Journal of Medical Science*. – 2022. – Т. 5. – С. 1-6.
10. Abdukodirov E. I. et al. Study of bioelectric activity of the brain in patients with neurosensorius deafness // *Oriental Journal of Medicine and Pharmacology*. – 2022. – Т. 2. – №. 05. – С. 10-19.
11. Karimovna X. F. Aspects of Anemia Related to Factors of the Lymphatic System // *International journal of health systems and medical sciences*. – 2022. – Т. 1. – №. 5. – С. 199-202.
12. Sulonovna E. Y. Peculiarities of the Development of Rickets in Children // *International journal of health systems and medical sciences*. – 2022. – Т. 1. – №. 5. – С. 159-163.
13. Абдуллаев Ш. Ю., Архипова М. Х. Использование новых биологически совместимых материалов при восполнении дефектов челюсти // *Стоматология*. – 1999. – Т. 78. – №. 3. – С. 37-38.
14. Абдуллаев Ш., Юсупова Д., Раимкулова Д. Значение сосудистого фактора в процессе заживления послеоперационных рубцов лица // *in Library*. – 2022. – Т. 22. – №. 4. – С. 125-127.
15. Абдуллаев Ш., Халилов А., Алимжанов Х. Применение обогащенной тромбоцитами плазмы в хирургической стоматологии // *Медицина и инновации*. – 2021. – Т. 1. – №. 2. – С. 80-85.

16. Yu, Abdullayev Sh, and N. Yu Muratova. "Application of osteoplastic compositions in mandibular endoprosthetics." *Conferencea* (2022): 263-264.
17. Gaffarova V. F. et al. Clinic-ecg Correlation Somatogenous of Conditioned Febrile Seizures in Children //International Journal of Human Computing Studies. – 2021. – Т. 3. – №. 1. – С. 114-116.
18. Isroilovich A. E., Kodirovich K. N., Jumanazarovich M. R. Hereditary Diseases of the Nervous System, Their Prevalence and Epidemiological Status //Central Asian Journal of Medical and Natural Science. – 2022. – Т. 3. – №. 6. – С. 127-132.
19. Muratova, N. Yu, I. I. Khasanov, and Sh Yusupov. "Application of ultrasonic cavitation in treatment of the purification of wounds of the maximum-face region." (2017).
20. Юсупова, Д., Халилов, А., Жураев, Б., & Мухамедова, Ш. (2021). Лечение и профилактика послеоперационных рубцов лица на современном этапе. *in Library*, 21(1), 48-53.
21. Юсупова Д., Абдуллаев Ш., Халилов А. Prevention of the formation of postoperative hypertrophic scars on the face //in Library. – 2020. – Т. 20. – №. 4. – С. 24-26.
22. Муратова Н. Ю., Хасанов И. И., Юсупов Ш. Ш. Застосування ультразвукової кавітації для лікування гнійних ран щелепно-лицевої ділянки.
23. Abdukodirov E. I., Khalimova K. M., Matmurodov R. J. Hereditary-Genealogical Features of Parkinson's Disease and Their Early Detection of the Disease //International Journal of Health Sciences. – №. I. – С. 4138-4144.
24. Ganiev A. A., Abdullaev S. Y., Abdurahmonov S. Z. Combined treatment for early-stage skin cancer of the head and neck area //World Bulletin of Public Health. – 2021. – Т. 4. – С. 3-6.
25. Шокиров Ш. Т., Ганиев А. А., Зайнутдинов М. О. Остеопороз В Дентальной Имплантологии. Принципы лечения и профилактики (обзор) //Журнал теоретической и клинической медицины. – 2016. – №. 4. – С. 160-163.
26. Ogli, Abdukodirov Eldor Tolibjon, and Yuldasheva Risolatkhon Kobil Kizi. "eco composite materials using basalt rocks." *Проблемы современной науки и образования* 4 (173) (2022): 111-114.
27. Абдуллаев, Ш., Халилов, А., Адылов, З., & Алимжанов, К. (2021). Лечение осложнений, возникающих при переломах нижней челюсти. *Медицина и инновации*, 1(4), 140-148.
28. Imomjonovich I. I., Amirkulovna A. G. Current immunological problems in kidney transplantation //Web of Scientist: International Scientific Research Journal. – 2021. – Т. 2. – №. 09. – С. 24-28.
29. Imomjonovich I. I., Ulyanovna S. G. Immunogenetic changes in kidney transplantation //International Engineering Journal For Research & Development. – 2021. – Т. 6. – №. 3. – С. 3-3.