



## Planting Cauliflower Seeds in the Open Field

Pirimqulova Muxabbat Xayitovna <sup>1</sup>, Mamarajabov Samandarbek Faxriddinovich <sup>2</sup>,  
Abdullayev Muxtorjon Bobir o'g'li <sup>3</sup>, Ochilov Baxtiyor Ziyodulla o'g'li <sup>4</sup>

<sup>1, 2, 3, 4</sup> Termiz Institute of Agrotechnologies and Innovative Development

**Abstract:** moisture cauliflower is very good as the closest relatives of culture to see. The soil should be at least 70 percent of optimum moisture, does not let to dry. This is especially important during the growth of seedlings. It quickly began to flourish even when the lightweight construction of the soil is shallow or loss of productivity can lead to the formation of your head. Very important and according to the temperature.

**Keywords:** Peat tablets, tightly in the soil, seedlings, greenhouse, and superphosphate, and potassium sulfate, kostryulkalarda.

Open doublesrga gulkaram seeds emen.

Cauliflower prepared in advance of planting bed is prepared and agricultural techniques you need to follow certain rules. In the fall, after harvest of the main crop, the soil will begin to prepare for spring planting. Do thischun, weeds and insects necessary to remove it. About two weeks later is applied fertilizer - organic fertilizer is spending to 5 kg and 30 g of potassium and phosphorus square meters. Urug'lantirilgandan bed after a bayonet wrapped and left to rest until the spring has been. With the onset of the soil is the removal of heat and urea - 20 g per square meter. Note: dense planting cauliflower in the soil, that is, re-extracted it you do not need. In the fall you if you have the time to dig up the earth, you need to do it in the spring, but then compressed before planting and ythe way, you will need to fill light. Start on the quality of this effect: the need to comply with this rule, they are dense and should be delicious.

Cauliflower is usually carried out through planting the seedling. However, today, in the ground, planting the seeds for gardeners often apply. The beginning've matured yhappen fast enough in and planting the seeds for a meeting during the whole summer to get 20-day break can be done 3-4 times do not forget that. Most inearly seedlings for planting in greenhouses and greenhouses ingrown crops in the summer and autumn falls directly on the ground. When the air temperature falls, the seedlings with the film shielded don't forget. There are many varieties in the market today. To grow plants in a greenhouse matures early which is perfect. Recently cabbage is very good, only in the south have been grown and there are everywhere today. However, also in the greenhouse, the weather conditions very cool, can start as usual grow smaller. Makes the best cabbage I reach in the middle of the season. Get the crop seedlings in the early planting and planting in the garden allows you - as well as second and third. Sa varieties types usually lateripening in climatic conditions to lqin can't find the time.

Planting the seeds

The first task which is the perfect combination for the early summer cauliflower salad crop ingrown. Fall seedlings is carried out according to the following rules. The friendly and healthy seedlings, you should choose to get the only large seeds. Before planting, take the following steps: seeds wrapped in cloth and dipped into the hot water. Do not use it to boil your best option is to level 50. The seeds

were cooled in cold water is now activated and is namlab for 8 hours with a solution of potassium permanganate.

This drug is a perfect germination will give. Special boxes sprinkled prepared for planting the seeds. The soil qan every day, but fertilizer and recommended a mixture of humus best himself. In addition, 15 grams of superphosphate and potassium sulfate soil mixture should be added to the bucket. Cauliflower planting seeds almost to an open place. A little hydrated and in compacted soildepth of 0,5 mm will remove the seeds. They are covered with the same mixture and ythe light inthe land is covered with.

The optimum temperature for the cabbage +20 degrees. In such conditions the seedlings 4-5 days within will appear. This yposes the main risk here: the cabbage goes quickly stretched out, wolves appear similar after the temperature of the air +6 to the level falls. After about 5 days, it rises to 15 degrees - that is personalized to come. The easiest method to fulfill this condition - is your kostryulkalarda your seedlings, but the place to open a place for maneivr seed planting cabbage leaves. Covered greenhouse or edge is removed from the film, which opened. Thus, the temperature is regulated.



#### The seedlingscultivation of the world

If you're dealing with your seedlings planted, then you 8-10 days ago, you should choose one of seedlings. The process of the soil a bit before you should soak it in. Keep in mind that seedlings of excess moisture in the tank disorders the get go. After you have out, the trees should ovqatlantirilishi. The preparation fi 45-50 days lasts. Get ready for at least 5 true leaves, planting seeds and should have well-developed root system. Qattiqlashtiradigan seedling, sun, frost , and make sure to change

the temperature. To do this, before planting, the seedlings start out at the balcony or open out to the street. The last few days, the plants in the greenhouse during worthwhile to be fully determined. The night is cold; the film will not be removed.

Plant cultivation method: suitable for areas which

A primary for planting in open areas cabbage garlic should be noted that the vitreous has an advantage. Reduces the profitability of the seedlings raised and the quality of the product and reduces the risk you'll notice you left no: plants should be in accordance with their biological development hours.

Best dry seed planting cauliflower, very suitable for the southern region. But with the advent of early ripening varieties, this method is used by the siberian city up to all the gardeners. In this case, if the root system is less tarvaqalangan, will penetrate deep into the soil. Before planting, hybrid, or according to different characteristics, as well as the fertility of the soil, depending on labeling rows and 45-60 cm direct on the bed is carried out. When the first true leaves fidel, he should be lubricated. Remains a depth of 15 cm between plants, the seedlings china, we have six leaves, releasing when the dissolution process is repeated and again in the range of 15 cm between the plants should remain. other place.

Growing cauliflower feature

The amount of healthy vegetables and depends on the quality of her care. Weak to take care of the flower for the white should be the same. It need watering, feeding, pasture, hilling and should loosen the soil. Season diseases and should implement preventive measures against pests.

Regular watering should be. Earlier seedlings after planting cabbage in the week two times were watering. Less watering the plants can be planted. If you are the rain, then watering is carried out after the drying of the soil. Regular water and heavy rains with garden vegetables is not recommended. Their temperature is low if the temperature is low, this can cause the development of fungal diseases. Flowers density cauliflower and whoever experiences that keep moisture for them to hit can be covered with two or three leaves.

From planting plants in open areas, three weeks later, they need **nutrition with a solution of Mullein** each hole in the bottom of a liter and a half of bulk. 10 liters of water and ½ liter of solution, the composition of the preparation. Again after ten days, each plant is fed with a liter of solution, is added to drug him qoshiqni. **Crystals**. Times for every season vegetables during the bush sixto eight quarts with mineral fertilizers need feeding. You can use two liters of water was diluted in ten nitrofoskadan you qoshiqni.

I too like cauliflower various pests, adverse conditions, shows the effects of fungal diseases. Therefore, in the cultivation of various vegetables should perform preventive therapy. At the same time is not used chemical from the drug.

Planting seedlings into the open ground - the scheme and agricultural technology

Early ripening varieties sowed in the open ground and **the middle of April-in the middle**. The start of planting a week ago seedlings superphosphate (3 g), potassium chloride diluted in a liter of water (3 g) with fed. This increases the cold resistance of the century. In addition, the plant is a new living environment to teach, to shake should you require them.

Early varieties in the beginning of May - early may be planted in the prepared soil before planting the ground in mumkinkuzda planted. Each well in compost (humus), wood ash, superphosphate, urea (1 bucket, 2 cups, 2 tbsp .) make a mixture. The fertile soil is mixed with all the parts.

Basic requirements and conditions

Cabbage, which requires the light of the sun the temperature bir cauliflower cultivation of fruit ripening and ripening to achieve technical need for a different type at least 120-160 looking for a warm day. The group has quite enough to do this in the middle of the summer period. However, the growth of the complexity of this type of cabbage, it is very sensitive to temperature change, for the



reason that cold bed at a depth soylash too much power for the landing place from the sun and protection is required.

Cauliflower is required in the light of the sun. Good, dense, starting only open to grow in illuminated area at a high level in this case are connected. Long garden crops is too close, can have a negative impact on the quality of the product. The smell of cabbage the best: potatoes, cucumbers, tomatoes, beets, peas and other baklagiller. Radishes and turnips, radishes and cabbage as other types of "appropriate" ekma crop it, then pathogenic bacteria and fungus can remain in the soil of the sport.

Growing cabbage for the soil, heavy clay soil is not suitable or weak. These plants are good moisture and grown in sandy soil allows air garden filled with harvested. When using the beds of organic fertilizer to improve the nutritional value of cabbage is better: mullein or rotten manure, humus, compost mass, acidic peat.

Planting cauliflower, regular watering, especially in the initial stages of moisture should be planting. At the same time, the development of fungal diseases usually turg'unligi of water in the soil is the main cause, therefore, the frequency of irrigation and the amount of rain depending on weather conditions and the size is determined. As a rule, in the second half of the summer, the ripening of the crop when it happens, cabbage now non-irrigated. This is a very new technology that allows you to get a very strong plants. After taking tablets buy peat, just put in a plastic container and should be poured with warm water. After a while they will swell and the feed mixture is converted. In one or two seeds in each glass will fall.

Namlantirish from time to time to take care of the little seedlings with the soil and care when needed is to make. Of cabbage are planted in the greenhouse in approximately the same way. It "Svetlitsa" like filling with a special film is good. The light of the sun take the best of such material and higroskopik (condensate yig'ilmaydi).

Also, you can grow cauliflower heads basements or cellars. In this case, the bound plants and root hung on the fence. The end of the growth process mass per 0 cabbage, 5 kg ha, while 15-20 smga in diameter will.

2-3 true leaves of plants with the formation of ammonium 5 g and 15 g of potassium nitrate was diluted with a bucket of water, with a solution of potassium or ammonium nitrate was feeding. 3-4 leaves are harvested when the plant is in the stomach, feeding from boric acid and molybdenum acid ammonium solution (10 liters of water for 2 g) by spray is carried out. Rarely were watering seedlings, but for many, the amount havalandırılır the room after irrigation. Open from planting to the place about two weeks ago, the shades of the seedlings to the temperature of the outside environment into the habit of plants, gradually hardened. 45-50 days at planting seedlings ready for planting. By this time, the plants should be at least 4-5 leaf.

Organic cabbage mixture will create favorable conditions for the development of the root system and all the necessary food, mis the delivery of oddalar. 1 meter from 4 kg to 8 kg organic matter and mineral fertilizers you need to produce 80-100 grams.

This is one of the products most recommended food. This has valuable medicinal properties and is very easy to digest. This is the best product to add to the baby first additional dishes.

Softened the soil before planting the seeds to a depth of 15-20 cm, the soil thoroughly rinse the seeds finely prepared cell was planted in the way. Rows in a range, and is similar to the distance when the distance between soketlar seedling planting. Sprinkle each hive 3-5 seeds, peat and sprinkled with a mixture of humus. The third true leaf formation in the plant along with two second left at home of crops, plants, and has become one of the most comfortable. Then planting care about, as well as in the case of the growth of the seedlings.

Wrapped box to the window from the window, a day later, sprinkled with the water of the soil. At a temperature of 18-20 degrees for 5-7 days during will appear. After the seedlings appear, its temperature was increased from 7-8 degrees to transplant the seedling to the room on will be. If this fails you, the seedlings quickly stretched out, is going to die.

Open planted into the ground.

About a week before planting the expected time held the feeding should be stopped. The only thing that can be useful - potassium chloride and superphosphate (3 g for 1 g water) solution add. This measurement, together with seedling hardening (at the time grow gradually go to a place where) will help to show resistance to fall, which can be the temperature.

Soil requirements and preparation

Neutral pH (6.7-7.5) the soil is preferred. Before any transfer seedling mineral complex with 1 g in the amount of insemination should 90-100 m<sup>2</sup>. Acidic soils "soil" or other deoksidlovchi stir. The extraction of the depth of the block is the shovel in the winter period (also, it deoksidlash) and on the basis of abnormal only in the spring season was in accordance with the purpose of getting laid off. Not necessarily a sufficient density of soil is important, it should not be too soft, otherwise the start is the same.

*The cauliflower seedless grow in the open ground*

This is a reliable method in the southern region, you can use the seeds is similar to planting seedlings, but immediately F. In cold areas, it is recommended to grow seedlings in the greenhouse or home. But you already warm or compost-heated seeds of the early varieties and the planting bed in the garden you can immediately. In this case the watch at a temperature of day and night, half a plastic bottle (taking part of the cap) to each of them a mini-greenhouse for the construction, deepening of the earth should.

Irrigation, soften.

Regular moisture and should be sufficient. Simple weather conditions in the week two times (until a power plant into) and then requires watering once (considering natural rainfall). Suddenly the bed for a bucket of water should be less than 1 square meter initially, but with the growth of consumption increased the level of your head. Watering of the earth between the flag and the flag can be combined with. Light and painless, rain the next day, remove weeds. Watering more frequently, the amount or the shredding of fungal infections and can lead to the development. It has a shallow root system, we should not forget that the vegetables are very weak and, therefore gevşemeyi through mulching also, to put instead of spreading too will be good.

Cauliflower varieties and hybrids.

Now, many varieties of cabbage, and there are hybrids. Too early, too late varieties. In addition, the color of exotic vegetables can be, for example, or orange salad.

Early ripening of the culture of short-term plant is able to the season. The beginning is very dense and surrounded with leaves. From mid-or late ripening cabbage leaves a little bit shortened. Ripening mid-ripening varieties of leaves out of rozetali at length the difference between. In addition, the head is at +20 ° c can be at a temperature of the formation. Early varieties and hybrids for the plant in the period of approximately 80-110 days.



Gregory F1 P3

A hybrid high-mahsul are included in the state register of the Russian federation. The beginning of your head to be able unusual shape: a mass of approximately 3 pounds.

Of cauliflower fresh and processed form, it is recommended to work in.

Cheddar F1

Ripe early spring and karotendan if the difference is in the perfect taste.

Cauliflower average weight 1 2 pounds. It is recommended to freeze for the winter, the beginning of cheddar f1.

**Conclusion:** the soil before planting the seeds to a depth of 15-20 cm softened, rinse thoroughly with the soil, the seeds finely prepared cell was planted in the way. Rows in a range, and is similar to the distance when the distance between soketlar seedling planting. Sprinkle each hive 3-5 seeds, peat and sprinkled with a mixture of humus. The third true leaf formation in the plant along with two second left at home of crops, plants, and has become one of the most comfortable. Then planting care about, as well as in the case of the growth of the seedlings.

## References

1. Khaitovna PM, Faksriddinovich MS Technology of growing cauliflower // Texas Journal of Interdisciplinary Research. - 2022. - T. 6. - S. 8-10.
2. [https://scholar.google.com/citations?view\\_op=view\\_citation&hl=ru&user=Buztwd0AAAAJ&autouser=1&citation\\_for\\_view=Buztwd0AAAAJ:u-x6o8ySG0sC](https://scholar.google.com/citations?view_op=view_citation&hl=ru&user=Buztwd0AAAAJ&autouser=1&citation_for_view=Buztwd0AAAAJ:u-x6o8ySG0sC)
3. [https://scholar.google.com/citations?view\\_op=view\\_citation&hl=ru&user=Buztwd0AAAAJ&autouser=1&citation\\_for\\_view=Buztwd0AAAAJ:2osOgNQ5qMEC](https://scholar.google.com/citations?view_op=view_citation&hl=ru&user=Buztwd0AAAAJ&autouser=1&citation_for_view=Buztwd0AAAAJ:2osOgNQ5qMEC)
4. [https://scholar.google.com/citations?view\\_op=view\\_citation&hl=ru&user=Buztwd0AAAAJ&autouser=1&citation\\_for\\_view=Buztwd0AAAAJ:qjMakFHDy7sC](https://scholar.google.com/citations?view_op=view_citation&hl=ru&user=Buztwd0AAAAJ&autouser=1&citation_for_view=Buztwd0AAAAJ:qjMakFHDy7sC)
5. <https://doi.org/10.5281/zenodo.7115076>
6. <https://doi.org/10.5281/zenodo.7131539>

7. Xayitovna, P. M., & Faxriddinovich, M. S. (2022). Types of corn grown in Uzbekistan and their peculiarities. *Texas Journal of Agriculture and Biological Sciences*, 3, 59-63.
8. Xayitovna, P. M., & Faxriddinovich, M. S. (2022). Cauliflower Growing Technology. *Texas Journal of Multidisciplinary Studies*, 6, 8-10.
9. Abdullayev, M., & Mamarajabov, S. (2022). VARIETIES AND ACHIEVEMENTS OF WHEAT PLANT SELECTION IN UZBEKISTAN. *Eurasian Journal of Academic Research*, 2(11), 100-104.
10. Khayitovna, P. M., & Faxriddinovich, M. S. (2022). PECULIARITIES OF GROWING CAULIFLOWER. *Science and innovation*, 1(D3), 144-146. Константинович, А. В., & Маслов, В. А. (2012). Выращивайте рассаду цветной капусты правильно. *Картофель и овощи*, (2), 25-26